Accepted Manuscript

Association between the Big Five factors of personality and survival seven years after heart transplantation

Roberto Sánchez-González, Amanda Rodríguez-Urrutia, Eila Monteagudo-Gimeno, Sandra Herranz-Villanueva, Félix Pérez-Villa, Luis Pintor-Pérez

PII: S0163-8343(19)30048-9
DOI: <https://doi.org/10.1016/j.genhosppsych.2019.02.005>
Reference: GHP 7395
To appear in: *General Hospital Psychiatry*
Received date: 11 February 2019
Revised date: 17 February 2019
Accepted date: 20 February 2019

Please cite this article as: R. Sánchez-González, A. Rodríguez-Urrutia, E. Monteagudo-Gimeno, et al., Association between the Big Five factors of personality and survival seven years after heart transplantation, *General Hospital Psychiatry*, <https://doi.org/10.1016/j.genhosppsych.2019.02.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Association between the Big Five factors of personality and survival seven years after heart transplantation

Authors:

Roberto Sánchez-González, MD, PhD ^{1,2,3}, Amanda Rodríguez-Urrutia, MD, PhD ⁴, Eila Monteagudo-Gimeno, MD ^{1,2,3}, Sandra Herranz-Villanueva, BSN ⁵, Félix Pérez-Villa, MD, PhD ⁶, Luis Pintor-Pérez, MD, PhD ⁵

¹ Department of Psychiatry. Institut de Neuropsiquiatria i Addiccions, Centre Emili Mira, Parc de Salut Mar. Barcelona. Spain.

² IMIM (Hospital del Mar Medical Research Institute). Barcelona. Spain.

³ Department of Psychiatry and Forensic Medicine. School of Medicine. Universitat Autònoma de Barcelona. Bellaterra (Cerdanyola del Vallès). Spain.

⁴ Consultation-Liaison Psychiatry Unit. Department of Psychiatry. Hospital Universitari Vall d'Hebron. Departament de Psiquiatria i Medicina Legal, Universitat Autònoma de Barcelona. Group of Psychiatry, Mental Health and Addiction, Vall d'Hebron Research Institute (VHIR). Barcelona. Spain.

⁵ Consultation-Liaison Service. Department of Psychiatry. Institut de Neurociències. Hospital Clínic i Provincial de Barcelona. Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS) – Universitat de Barcelona. CERCA Programme/Generalitat de Catalunya. Barcelona. Spain.

⁶ Heart Transplantation Division. Department of Cardiology. Thorax Institute. Hospital Clínic i Provincial de Barcelona. Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS) – Universitat de Barcelona. Barcelona. Spain.

Corresponding author:

Roberto Sánchez-González

Department of Psychiatry

Institut de Neuropsiquiatria i Addiccions, Centre Emili Mira, Parc de Salut Mar
C/ Prat de la Riba, 171, 08921, Santa Coloma de Gramenet, Barcelona (Spain)

Tel.: +34934628900

Fax: +34934683742

E-mail address: rsanchezgonzalez@psmar.cat

Financial disclosure:

The authors have no conflicts of interest or financial disclosures to report.

Article data:

Word length of the article's text: 776 words

Number of tables: 1

Number of figures: 1 (supplementary data)

Running title:

Personality and survival in heart transplantation recipients

Association between the Big Five factors of personality and survival seven years after heart transplantation

Keywords: Five-Factor Model, heart transplantation, personality, positive psychology, psychosocial, survival.

INTRODUCTION

Over the past decades, many studies have tried to search psychosocial predictors of survival after heart transplantation (HT), independent of the recipient's cardiac characteristics and risk factors [1-4]. Although some personality traits are well-known predictors of posttransplant clinical outcomes, the lack of empirical findings considerably limits the evidence about its real impact on mortality [5]. The objective of this study was to perform an exploratory analysis to investigate the relationship between the Big Five factors of personality and long-term survival in HT recipients.

METHODS

We performed a prospective study that included consecutive adult candidates who underwent HT at the University Clinical Hospital of Barcelona from January 2006 to December 2012. HT recipients were followed-up until May 2018 (maximum observation period of 12 years). The study protocol was approved by the Ethical Committee Board of the hospital. Prior to the surgery, we applied the Structured Clinical Interview for DSM-IV axis I and axis II disorders, Clinician Version (SCID), and the Hospital Anxiety and Depression Scale (HADS) for the evaluation of psychopathology. The Apgar-Family questionnaire was used to determine family function. For the personality assessment, we used the Spanish validation of the Five Factors Inventory Revised (NEO-FFI-R) [6]. It is a self-report questionnaire with 60 items that measures the Big Five main personality factors: Neuroticism, Extraversion, Openness to experience, Agreeableness and Conscientiousness. Since there are no generally accepted cut-off points for NEO-FFI-R, we dichotomized each personality factor using the mean scores obtained in the Spanish validation [6]. For the statistical methods used to evaluate each personality factor as a predictor for post-HT survival, we conducted the Cox proportional hazard regression analysis to test multivariate associations with time until death. This analysis was adjusted to consider the main medical and psychosocial variables. The Kaplan–Meier survival analysis was performed for each dichotomized factor separately, comparing the HT recipients with either low or high scores and using the Log-Rank (Mantel-Cox) chi-square test to check differences between survival curves.

RESULTS

We included 78 adult HT recipients in the follow-up study protocol. The baseline characteristics of this sample were published in a previous paper [7]. The mean follow-up after HT was 7 years (SD = 3.52 and range 0 – 12.33 years), and at the end of this period, 53 recipients were alive (68%) and 25 had died (32%). The mean scores of each personality factor were: 20.47 (SD = 7.66) for Neuroticism, 28.06 (SD = 6.20) for Extraversion, 29.53 (SD = 7.07) for Openness to Experience, 34.28 (SD = 6.04) for Agreeableness and 35.02 (SD = 6.19) for Conscientiousness. In the multivariate analysis, only Agreeableness was independently associated with survival (Table 1). This result was consistent with the Kaplan–Meier analysis (Figure 1, supplementary data)

that showed that lower scores in Agreeableness were significantly associated with an increased risk of death, as indicated by the results of the Log-Rank chi-square test: $\chi^2 = 4.83$, $df = 1$, $p = 0.029$.

DISCUSSION

This is the first study to investigate the relation between the Big Five factors of personality and long-term survival in HT recipients. Our results suggest that higher levels of Agreeableness may be independently associated with better clinical outcomes and post-HT survival. Previous investigations that are in accordance with our results, demonstrated a relationship between high Agreeableness and both a lower level of clinical disorders and higher optimism [8,9]. Likewise, there is growing evidence linking positive psychological characteristics, like optimism and resilience, to better health outcomes, and in the last years, optimism-specific interventions for people with cardiovascular disease have been developed [10]. In contrast with the results obtained by Stilley et al. (2004), which showed a prevalence of 33% of personality disorders among cardiothoracic transplant recipients according to the Five-Factor model, the mean scores of NEO-FFI-R of our sample were similar to those of the general population [5]. Regarding the other four personality factors, the typical pattern linked to clinical disorders are high Neuroticism, low Conscientiousness, and low Extraversion, but these factors were not associated with long-term survival in our study [8]. This finding may be explained by the exhaustive pre-transplant selection criteria, which might generate personality traits among our sample that are highly homogenous. The main limitations of our study are that we have performed an exploratory analysis based in only 25 deaths, there are multiple comparisons and a more conservative analysis would use $p = 0.01$ as significant. Based on our results, we postulate that adaptive personality traits such Agreeableness could establish a construct opposite of Type A behavior or a vulnerability to disease model, play a role as a protective factor and a long-term survival predictor in HT recipients, and provide a valuable target for new therapeutic strategies.

REFERENCES

- [1] Sánchez R, Baillés E, Peri JM, Bastidas A, Pérez-Villa F, Bulbena A, Pintor L. Cross-sectional psychosocial evaluation of heart transplantation candidates. *Gen Hosp Psychiatry*. 2014;36(6):680-5.
- [2] Dew MA, Rosenberger EM, Myaskovsky L, DiMartini AF, DeVito Dabbs AJ, Posluszny DM, et al. Depression and anxiety as risk factors for morbidity and mortality after organ transplantation: a systematic review and meta-analysis. *Transplantation*. 2015;100(5):988-1003.
- [3] Spaderna H, Zittermann A, Reichenspurner H, Ziegler C, Smits J, Weidner G. Role of Depression and Social Isolation at Time of Waitlisting for Survival 8 Years After Heart Transplantation. *J Am Heart Assoc*. 2017 Nov 29;6(12).
- [4] Farmer SA, Grady KL, Wang E, McGee EC Jr, Cotts WG, McCarthy PM. Demographic, psychosocial, and behavioral factors associated with survival after heart transplantation. *Ann Thorac Surg*. 2013;95(3):876-83.

- [5] Stilley CS, Dew MA, Pilkonis P, Bender A, McNulty M, Christensen A, et al. Personality characteristics among cardiothoracic transplant recipients. *Gen Hosp Psychiatry*. 2005;27(2):113-8.
- [6] Aluja A, García O, Rossier J, García LF. Comparison of the NEO-FFI, the NEO-FFI-R and an alternative short version of the NEO-PI-R (NEO-60) in Swiss and Spanish samples. *Pers Individ Dif*. 2005;38(3):591-604.
- [7] Sánchez R, Baillés E, Peri JM, Bastidas A, Pérez-Villa F, Bulbena A, Pintor L. Assessment of psychosocial factors and predictors of psychopathology in a sample of heart transplantation recipients: A prospective 12-month follow-up. *Gen Hosp Psychiatry*. 2016;38:59-64.
- [8] Malouff JM, Thorsteinsson EB, Schutte NS. The relationship between the five-factor model of personality and symptoms of clinical disorders: A meta-analysis. *J Psychopathol Behav Assess*. 2005;27(2):101-14.
- [9] Sharpe JP, Martin NR, Roth KA. Optimism and the Big Five factors of personality: Beyond Neuroticism and Extraversion. *Pers Individ Dif*. 2011;51(8):946-51.
- [10] Mohammadi N, Aghayousefi A, Nikrahan GR, Adams CN, Alipour A, Sadeghi M, et al. A randomized trial of an optimism training intervention in patients with heart disease. *Gen Hosp Psychiatry*. 2018;51:46-53.

	Neuroticism		Extraversion		Openness to experience		Agreeableness		Conscientiousness	
	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value
Main factor *	0.33 (0.04 - 2.89)	0.316	1.12 (0.4 - 4.16)	0.829	0.45 (0.08 - 2.43)	0.355	0.11 (0.01 - 0.71)	0.021	0.47 (0.12 - 1.85)	0.282
Illness duration before HT (years)	0.86 (0.76 - 0.97)	0.016	0.79 (0.67 - 0.93)	0.004	0.85 (0.75 - 0.95)	0.006	0.82 (0.72 - 0.94)	0.004	0.85 (0.76 - 0.96)	0.009
Male sex	1.5 (0.34 - 6.61)	0.592	0.94 (0.2 - 4.39)	0.904	1.63 (0.37 - 7.26)	0.519	1.91 (0.43 - 8.52)	0.396	1.19 (0.3 - 4.72)	0.809
Ejection fraction (%)	0.99 (0.93 - 1.05)	0.709	0.99 (0.92 - 1.06)	0.765	0.98 (0.93 - 1.05)	0.633	0.99 (0.92 - 1.05)	0.671	0.98 (0.91 - 1.04)	0.777
Pulmonary Vascular Resistance (UW)	1.19 (0.6 - 2.36)	0.626	1.36 (0.71 - 2.6)	0.356	1.32 (0.6 - 2.9)	0.486	1.41 (0.76 - 2.58)	0.273	1.14 (0.59 - 2.2)	0.866
Pulmonary Artery Pressure (mmHg)	1 (0.93 - 1.07)	0.997	0.99 (0.93 - 1.07)	0.953	0.99 (0.92 - 1.07)	0.833	0.99 (0.92 - 1.07)	0.866	1 (0.94 - 1.07)	0.924
Creatinine (mg/dL)	1.57 (0.75 - 3.25)	0.228	2.02 (0.85 - 4.79)	0.112	1.51 (0.74 - 3.07)	0.257	2.54 (0.98 - 6.59)	0.054	1.41 (0.69 - 2.88)	0.347
AST/ALT ratio	2.53 (1.07 - 5.95)	0.034	2.18 (1.03 - 4.6)	0.042	1.96 (0.91 - 4.23)	0.088	3.44 (1.35 - 8.78)	0.001	2.01 (0.97 - 4.16)	0.066
DSM-IV diagnosis	1.12 (0.24 - 5.26)	0.888	1.17 (0.26 - 5.21)	0.837	1.36 (0.27 - 6.87)	0.706	1.81 (0.39 - 8.45)	0.453	1.09 (0.22 - 5.27)	0.915
HADS total score	1.03 (0.21 - 5.16)	0.971	0.26 (0.05 - 1.35)	0.109	0.43 (0.08 - 2.37)	0.303	0.49 (0.11 - 2.19)	0.533	0.54 (0.12 - 2.42)	0.419
Apgar-Family questionnaire score	1.62 (0.35 - 7.56)	0.539	1.5 (0.33 - 6.79)	0.606	1.30 (0.3 - 5.67)	0.726	1.03 (0.23 - 4.54)	0.965	1.33 (0.31 - 5.8)	0.702

Table 1. Cox Proportional-Hazard Ratios for pre-HT recipient characteristics, the Big Five factors of personality measured with NEO-FFI-R, and time until death (n = 78).

ALT = Alanine Transaminase; AST = Aspartate Transaminase; CI = Confidence Interval; DSM-IV = Diagnostic and Statistical Manual of Mental Disorders, 4th Edition; HADS = Hospital Anxiety and Depression Scale; HR = Hazard Ratio; HT = Heart Transplantation.

* Main factor refers to Neuroticism, Extraversion, Openness to experience, Agreeableness and Conscientiousness, respectively.

Neuroticism

Extraversion

Openness to Experience

Agreeableness

Conscientiousness

--- Over mean
— Below mean

Figure 1