Journal Pre-proof

Age at illness onset and physical activity are associated with cognitive impairment in patients with current diagnosis of Major Depressive Disorder

Yolanda Sánchez- Carro , Maria J. Portella , Itziar Leal-Leturia ,
Neus Salvat- Pujol , Mikel Etxandi , Aida de Arriba-Arnau ,
Mikel Urretavizcaya , Esther Pousa , Alba Toll , Pilar Álvarez ,
Virginia Soria , Pilar López-García

PII: S0165-0327(20)32876-7
DOI: <https://doi.org/10.1016/j.jad.2020.10.032>
Reference: JAD 12574

To appear in: *Journal of Affective Disorders*

Received date: 14 January 2020
Revised date: 4 September 2020
Accepted date: 12 October 2020

Please cite this article as: Yolanda Sánchez- Carro , Maria J. Portella , Itziar Leal-Leturia ,
Neus Salvat- Pujol , Mikel Etxandi , Aida de Arriba-Arnau , Mikel Urretavizcaya , Esther Pousa ,
Alba Toll , Pilar Álvarez , Virginia Soria , Pilar López-García , Age at illness onset and physical ac-
tivity are associated with cognitive impairment in patients with current diagnosis of Major Depressive
Disorder, *Journal of Affective Disorders* (2020), doi: <https://doi.org/10.1016/j.jad.2020.10.032>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V.

current diagnosis of Major Depressive Disorder

Age at illness onset and physical activity are associated with cognitive impairment in patients with current diagnosis of Major Depressive Disorder

Yolanda Sánchez- Carro^{a,b,c}, Maria J. Portella^{c,d}, Itziar Leal-Leturia^{a,b,c}, Neus Salvat-Pujol^e, Mikel Etxandi^e, Aida de Arriba-Arnau^e, Mikel Urretavizcaya^{e,c,f}, Esther Pousa^g, Alba Toll^h, Pilar Álvarez^{c,h}, Virginia Soria^{e,c,f}, Pilar López-García^{a,b,c}

^a Department of Psychiatry, Universidad Autónoma de Madrid, Spain.

^b Department of Psychiatry, Instituto de Investigación Sanitaria Princesa (IIS Princesa), Spain.

^c Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Carlos III Health Institute, Spain.

^d Biomedical Research Institute Sant Pau (IIB-Sant Pau), Universitat Autònoma de Barcelona (UAB), Catalonia, Spain.

^e Bellvitge University Hospital, Psychiatry Department. Bellvitge Biomedical Research Institute (IDIBELL), Neurosciences Group - Psychiatry and Mental Health. Barcelona, Spain.

^f Department of Clinical Sciences, School of Medicine, Universitat de Barcelona, Barcelona, Spain.

^g Servei de Psiquiatria, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain.

^h Institute of Neuropsychiatry and Addictions, Hospital del Mar, IMIM, Barcelona, Spain.

Corresponding author: Yolanda Sánchez Carro, Department of Psychiatry, School of Medicine. Universidad Autónoma de Madrid. 4 Arzobispo Morcillo Street - Madrid (Spain). ZIP CODE: 28029. Contact: yolanda.sanchezc@uam.es /+34658769786

Highlights

current diagnosis of Major Depressive Disorder

1. MDD patients show cognitive deficits with global and specific domain impairment.
2. Years of schooling, job status, physical activity and the age at illness onset might predict the global cognitive deficits in MDD patients.
3. Physical activity and reinforcement of cognitive reserve should be incorporated in the assessment and treatment protocols in MDD.

Abstract

Background: Cognitive impairment has been reported in patients with Major Depressive Disorder (MDD). This study aims to explore the association between lifestyle habits and health-related factors and the presence of cognitive symptoms in MDD patients.

Methods: Demographic, clinical, health-related variables and cognitive scores measured with the Cambridge Neuropsychological Test Automated Battery (CANTAB) were compared between 74 patients with current MDD and 68 healthy controls (HC). To test the hypothesis of associated factors to cognitive symptoms, multivariate backward stepwise linear regression models were run.

Results: Significant neuropsychological deficits were evident in MDD compared with HC in the global cognitive index ($F=8.29$; $df=1, 140$; $p=0.005$). In the regression analysis performed on MDD and HC, years of schooling ($\beta=-0.11$; $p<0.001$), job status ($\beta=-0.50$; $p=0.016$), physical activity ($\beta=-0.25$; $p=0.04$) and age at illness onset ($\beta=0.17$; $p=0.017$) were statistically significant factors associated to cognitive impairment. The regression model ran in HC showed that only years of schooling were significant ($\beta=-0.07$; $p<0.001$) in this group.

Limitations: Sample size was relatively small. Everyday cognitive skills were not evaluated.

Conclusions: MDD patients have cognitive deficits. These deficits are linked with the years of education, job status, age of onset of the disease and the performance of physical activity. These results support the importance of the implementation of interventions targeting the cognitive reserve and lifestyle habits of MDD patients, in addition to the conventional therapeutic approach focused on symptoms control.

Keywords

Major depressive disorder; cognitive impairment; years of schooling; health-related variables; physical activity; age of onset.

1. Introduction

Major Depressive Disorder (MDD) is a common psychiatric disorder and a leading cause of disability (World Health Organization (WHO), 2017). Cognitive impairment is a clinical feature that is present in most MDD patients and mainly affects memory, attention, psychomotor speed and executive function. These cognitive deficits are key contributors to functional disability and increased risk of depression relapse (Culpepper et al., 2015). Although diagnosis and treatment approaches for MDD are mainly focused on affective symptoms (Kennedy et al., 2002), cognitive deficits should be taken into account since they mediate functional impairment during symptomatic phases from the first episode (Lam et al., 2014) and also during periods of remission (Rock et al., 2014; Baune and Air, 2016).

Despite its clinical relevance, not all MDD patients show cognitive deficits. The evolution of the cognitive deficits along the course of the illness may vary, for this

reason is important to know the factors that could be associated to the presence of cognitive impairment in MDD. Evidence suggest that patients with midlife-onset and late-onset depression, treatment resistant depression and with a greater number of depressive episodes have worse cognitive performance, especially in executive functioning (Paelecke-Habermann et al., 2005; Zheng et al., 2015; Eraydin et al., 2019). However, cognitive symptoms have also been reported in medicated (Yoshida et al., 2012) and remitted patients, as a result of the functional and anatomical changes in their brain (Kanner, 2004; Jannssen et al., 2007; Salvat-Pujol et al., 2017), and in patients with a first episode of depression (Chen et al., 2015). Besides, it has been observed that cognitive domains such as verbal learning, memory, verbal fluency and psychomotor speed are more sensitive to the severity of the episode (Douglas and Porter, 2009; Davis et al., 2017) and it has been verified a great impairment of executive functions in patients with a late onset of the disease (Eraydin et al., 2019). Other lifestyle-related factors such as physical activity, tobacco use, etc could be associated with cognitive impairment in MDD, as it is the case in other psychiatric conditions (Depp et al., 2015; Aas et al., 2019). However, little is known about the effect of the interaction of the lifestyle-related factors on cognitive performance in MDD, as few studies have explored the relationship between lifestyles and cognitive symptoms (Breslau et al., 1993; Pareja-Galeano et al., 2016).

By contrast, there is some evidence about the benefits of physical exercise (as an example of healthy lifestyle) on cognitive performance in clinical and non-clinical populations (Kramer and Erickson, 2007; Cassilhas et al., 2016; Karssemeijer et al., 2017) although neurobiological mechanism behind the beneficial effects of physical

exercise are unclear. Current hypotheses accept that physical exercise induce both acute and chronic effects affecting several putative systems implicated in depression such as neuroendocrine system, neurogenesis, oxidative stress, auto-immunity and producing cortical structural changes (for a review see Schuch et al, 2016). Decreases in arousal (Lambourne et al., 2010), increased catecholamines release (McMorris et al., 2009) and brain derived neurotrophic factor (BDNF) levels (Ferris et al., 2007), changes in cerebral activation (Meusel et al, 2013; Schneider et al, 2019) and enhanced brain plasticity have been found after physical activity both in healthy individuals and depressive patients (Gujral et al., 2017), leading to improved performance in specific cognitive domains: attention control (Colcombe et al., 2004), spatial memory (Erickson et al., 2011) and executive functioning (Kubesch et al., 2003; Colcombe et al., 2003). This cognitive improvement is higher with moderate physical activity and when it is performed simultaneously to the cognitive task (Rathore and Lom, 2017).

Another lifestyle factor that may affect cognitive performance is tobacco use. Nicotine may have effects on motor abilities, attention and memory which may have implications for initiation of smoking and maintenance of tobacco dependence (Heishman et al., 2010). However, chronic smoking is associated with worse cognitive performance (Durazzo et al., 2010; Wagner et al., 2012). Tobacco use prevalence in mental health patients is two to three times higher than in general population (Lasser et al., 2000). MDD patients are more prone to smoke and have greater difficulty quitting smoking (Prochaska, 2011; Weinberger et al., 2017). Smoking status has not been associated with cognitive impairment in MDD (Morisano et al., 2013). The new research should include larger sample sizes and longitudinal designs, as well as

evaluate the effects of smoking abstinence and reinstatement on cognition. This type of studies may be relevant due to the prevalence of this habit in these patients.

All these lifestyle factors, impact on metabolism, and people with non-healthy life habits are more prone to develop a metabolic syndrome (MetS). MetS is a cluster of cardiovascular risk factors including abdominal obesity, hypertension, hyperglycemia, low high-density lipoprotein cholesterol (HDL-C) and elevated triglycerides (Cleeman et al., 2001). MetS has a higher incidence in patients with psychiatric conditions, such as schizophrenia, bipolar disorder and MDD (Mitchell et al., 2011; Vancampfort et al., 2013; Vancampfort et al., 2014). As in the rest of psychiatric disorders, the relation between MetS and MDD seems to be bidirectional. Both share underlying etiological mechanisms like chronic inflammation or insulin resistance as observed in cohort studies where depressed individuals showed greater risk of having MetS that contribute to the chronicity of MDD (Vogelzangs et al., 2014; Marijnissen et al., 2017), and, patients with MetS showing higher comorbidity with MDD (Pan et al., 2012). It has been observed in animal models that the metabolic syndrome can cause cognitive problems, due to the correlation between the volume of white matter and HDL-C and a negative correlation between fasted blood glucose levels and hippocampal cell density (Livingston et al., 2020).

Studies in patients with diabetes have found evidence of poorer cognitive outcomes associated with the presence of depressive symptoms (Dana et al., 2016), with patients with type II diabetes having an increased risk of cognitive dysfunction (Sullivan et al., 2013).

However, the link between cognition and lifestyle seems to be bidirectional (Allan et al., 2014). Given that preserved executive functions are needed to plan, initiate, monitor and control actions, it has been postulated that executive functions facilitate the engagement in goal-directed health behaviours as physical activity, healthy diets and tobacco and drugs cessation. In this line, a strong reciprocal relationship between executive functions and physical exercise has been described in older adults (Daly et al., 2014).

Taken together, the relation of lifestyle with cognitive functioning in patients with MDD is far from being well understood. Therefore, the aim of the present study is to examine the association between depressive symptoms, lifestyle habits such as tobacco use and performance of physical exercise, and metabolic syndrome like a health-related factor and the presence of cognitive symptoms in MDD patients.

2. Methods

In this cross-sectional study the sample comprised 142 participants (74 MDD patients and 68 healthy controls (HC). All patients were recruited from primary healthcare centers of Madrid and the outpatients' psychiatric services of La Princesa University Hospital (Madrid), Bellvitge University Hospital and Hospital del Mar (Barcelona).

Patients had to fulfil criteria for a current episode of MDD according to the Diagnostic and Statistical Manual of Mental Disorder 4th Edition (DSM-IV-TR) (American Psychiatric Association, 2000), which was confirmed with a MINI interview. HC were recruited from the community and they were matched in terms of sex and age with patients, absence of current or past major psychopathology was discharged with the MINI.

current diagnosis of Major Depressive Disorder

Patients and HC had to meet the following inclusion criteria: 18-69 years old, Caucasian with European ancestry, ability to understand and to sign the informed consent. The exclusion criteria were: presence of any neurological or inflammatory disorder, use of antibiotic and/or anti-inflammatory treatment, and pregnancy (for all subjects); and absence of current or past mental disorders different to MDD, or treatment with electroconvulsive therapy (ECT) for the present episode and in the last 6 months (for patients). The study was carried out in accordance with the declaration of Helsinki and following the good practice guidelines. The Local Ethics Committees of each hospital approved the research protocol Clinical Research Ethics Committee (CEIC) of La Princesa University Hospital, CEIC of Bellvitge University Hospital and CEIC of the Parc de Salut Mar.

All participants provided written informed consent after having received a full explanation of the study.

2.1 Demographic, clinical and lifestyle variables

Demographic variables assessed in all participants were age, sex, marital status, estimated intelligence quotient (IQ) through the vocabulary subtest from the Wechsler Adult Intelligence Scale-Third Edition, years of education and current job status.

Clinical variables were obtained from the sample of MDD patients, including: age at illness onset, Hamilton Depression Rating Scale-17 items (HDRS-17) and stage of illness measured by the Maudsley Staging Method (MSM), which is a multidimensional staging model (Fekadu et al., 2009, Fekadu et al. 2018) that covers:

1. Duration of depressive episode, according to the following categories: acute (≤ 12 months), subacute (13 to 24 months) and chronic (> 24 months);

current diagnosis of Major Depressive Disorder

2. Severity of current episode measured by the Clinical Global Impression scale (CGI), that classifies the current episode of depression into remission, subthreshold, mild, moderate, severe and very severe patients and
3. Treatment failures (resistance). Thase and Rush scale was used to measure treatment resistance. Patient's depressive episode was classified with severity scores from 2 to 13 ranged as mild 2-5, moderate 6-9 and severe 10-13 (See the procedure in supplementary material).

Antidepressant medication (mg/day) consumed by patients was also recorded:

Selective Serotonin Reuptake Inhibitors (SSRI) such as Citalopram, Fluoxetine, Paroxetine, Sertraline, Fluvoxamine and Escitalopram, Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs): Venlafaxine and Duloxetine, Tricyclic

Antidepressants (TCAs): Amitriptyline and Imipramine and Tetracyclic antidepressants (TeCAs): Mirtazapine.

Other scales were administered to all participants: World Health Organization Disability Assessment Schedule 2.0. 12 items (WHODAS) that measures the functional state of the subject and State-Trait Anxiety Inventory (STAI: S-Anxiety and T-Anxiety) in order to measure the anxiety.

Lifestyles were explored through related variables such as tobacco use (subjects were classified as smokers, abstainers and non-smokers) and general physical activity measured using the International Physical Activity Questionnaire (IPAQ) by ranging "regular activity" and "last week activity" into low, moderate and high exercise. Other health-related variables and data were directly measured and collected at each center, for the present study at the time of the clinical assessment. Records included: body

current diagnosis of Major Depressive Disorder

mass index (BMI) (kg/m^2), waist circumference (cm), blood pressure (diastolic blood pressure and systolic blood pressures mm/Hg), fasting blood sugar level (mg/dL), HDL-C and triglycerides level (mg/dL). High values in these variables are known to be part of the MetS (Cleeman et al., 2001) (See in supplementary material).

2.2 Neuropsychological assessment

Cambridge Neuropsychological Test Automated Battery (CANTAB) was used to evaluate cognitive performance of all participants. This battery has been extensively utilized across neuropsychiatric patients (Levaux et al., 2007; Kim et al., 2014), and has been validated for major depression (Sweeney et al., 2000).

Five CANTAB tests were administered on a desktop computer with a touch-sensitive screen, and covered the following cognitive domains:

- Motor Screening Task (MTS): speed of processing.
- Delayed Matching to Sample (DMS): short-term memory.
- Spatial working memory (SWM): strategy use and working memory.
- One Touch Stockings of Cambridge (OTS): spatial planning (executive function).
- Rapid visual information process test (RVP): visual sustained attention.

2.3 Statistical Analysis

All data were analyzed using IBM SPSS Statistics (IBM SPSS Statistics for Windows, Version 24.0). Demographic, clinical, health-related and lifestyle variables were compared between HC and MDD patients by means of χ^2 test (or Fisher's exact test when expected frequencies in each cell were lower than 5) for categorical variables and t-tests for continuous variables. Cognitive raw scores were also compared between groups using ANOVA tests (p-value correction was applied to control for multiple testing).

Cognitive functioning was used as the primary variable of the present study; raw scores were z-transformed in a way that higher scores meant greater cognitive deficit –therefore some of the scorings were inverted– and were adjusted by age, IQ and gender using a regression model, before comparing HC and MDD patients. In order to obtain a global cognitive dysfunction index (GCDI) an exploratory factor analysis, using principal component analysis (PCA) to extraction method, was carried out with the whole sample, including all the tests from the CANTAB. Unrotated solution was then used as the dependent variable.

To test the hypothesis of associated factors to cognitive symptoms, different intermediate regression models were run in the sample of patients and HC. At first, a regression model was made with the scores obtained by the subjects in GCDI with the sociodemographic variables as predictors, observing that IQ, age and gender were part of a significant regression model in the prediction of the dependent variable. For this reason, in the final predictive model, the variables were adjusted by these three variables. A correlation study was performed with the Pearson test (GCDI) and years of schooling, age at illness onset (measured in years) and HDRS-17, WHODAS, STAI-S and STAI-T scores. Spearman test was performed to know the relationship with ordinal variables: tobacco use (smokers/nonsmokers/abstainers), physical activity (low/moderate/high) and severity of illness categorization by MSM (mild/moderate/severe). With Metabolic Syndrome variable (having or not metabolic syndrome) and job status (active or non-active status), a t-test was done. Due to the high number of clinical variables that correlate with the dependent variable, an intermediate regression analysis was performed to determine the non-explanatory variables and eliminate them from the final model.

After this, significantly associated variables were finally entered into one block of independent variables, and clinical variables of depression were entered into a second block of independent variables. By doing this, it was possible to contrast the variance percentage explained by the model in the sample of patients and controls by the variables, observing the change in the R^2 when introducing the predictive clinical variable. Once the model was carried out, compliance with the assumptions of non-multicollinearity was checked by checking the statistics of the Variance Inflation Factor

(VIF). The independence of the residuals was verified using the Durbin-Watson test, homoscedasticity and linearity were also verified using the dispersion graph of the standardized residuals against the predicted standardized scores. By means of the Kolmogorov-Smirnov test, compliance with the normal distribution of residues was contrasted.

Finally, to know the possible interactions between variables with the current MDD episode, a univariate linear regression was carried out.

3. Results

Both groups were comparable in terms of age, sex and civil status (Table 1). Patients completed fewer years of education and had a lower premorbid IQ and were more disabled and inactive than HC (Table 1). Patients had higher scores on state and trait anxiety. In addition, patients displayed worse psychosocial functioning, having difficulties in carrying out activities of daily living. In the clinical scales of severity of the disease, patients obtained scores suggestive of a moderate-severe episode (MSM $\bar{X}$ = 7.31, SD= 2.07; HDRS-17 $\bar{X}$ = 20.01, SD= 5.41).

Life-style habits and health-related variables were similar between the two groups, except for physical activity, since patients underwent physical activity less frequently than HC. The percentage of metabolic syndrome between HC and patients did not reach a significant difference (HC = 23. 1%; MDD = 16.2%).

PCA revealed a one-factor solution based on eigenvalues above 1, that was used as a score of global cognitive dysfunction for each participant. This factor represented the five domains scores: attention, memory, executive function, latency and impulsivity (see the loadings in Supplementary Material Table 3) and explained a 53.1% of variance with a partial correlation coefficient between variables Kaiser-Meyer-Olkin (KMO) = 0.786).

MDD patients showed worse global cognitive performance in cognitive tests from the CANTAB (omnibus $F=8.29$; $df=1, 140$; $p=0.005$); with poorer scores than HC in memory, latency of psychomotor speed and correct trials on executive function, considering p -value below 0.01 (Table 2).

Correlation and mean difference analyzes performed to explore the relationship between the different variables with the score obtained by the subjects on the GCDI revealed a relationship between this variable and the sociodemographic variables: years of education ($r= -0.58$, $p < 0.001$), job status ($t= 6.11$, $p < 0.01$), life-style factors: physical activity ($r=-0.19$, $p = 0.023$), and clinical variables: age at illness onset ($r= 0.43$, $p < 0.001$), consumption of SNRI ($\rho = 0.38$, $p < 0.001$) and scores on the WHODAS ($r= 0.23$, $p = 0.007$), STAI-T ($r= 0.24$, $p = 0.004$) and STAI-S ($r= 0.19$, $p = 0.025$) scales.

An intermediate regression model was built up with the different clinical variables: age at illness onset, consumption of SNRI and WHODAS, STAI-T and STAI-S scores, to know which one was the more relevant to introduce in the final regression model. Age of illness was the only variable with statistical significance among all the clinical variables that had had a significant correlation with the dependent variable ($\beta = 0.033$; $p < 0.001$).

In the sample of MDD patients the final model which predictors were years of schooling (measured in years), job status (two level: active/non-active), physical activity (low/moderate/high) and age at illness onset (measured in years) was statistically significant ($F=18.39$; $p < 0.001$) and explained 52.3% of the variance. The significant factors were years of education ($\beta=-0.11$; $p < 0.001$), job status ($\beta=-0.50$;

current diagnosis of Major Depressive Disorder

$p=0.016$), physical activity ($\beta=-0.25$; $p=0.04$) and age at illness onset ($\beta=0.17$; $p=0.017$).

All factors showed $VIF<10$ and tolerance >0.1 .

By studying the residuals, their independence ($DW= 1.9$), homoscedasticity, linearity and adjusted to normal distribution (K-S test $p = 0.20$) were confirmed. Figure 1 shows the scatterplot for the final model.

In the sample of HC, the final model which predictors were years of schooling, job status and physical activity was statistically significant too ($F=6.98$; $p<0.001$) and explained 24.9% of the variance. The significant factor was years of education ($\beta=-0.07$; $p<0.001$). Job status ($\beta=0.18$; $p=0.56$) and physical activity ($\beta=-0.09$; $p=0.52$) were not significant in the regression model. All factors showed $VIF<10$ and tolerance >0.1 .

By studying the residuals, their independence ($DW = 1.5$), homoscedasticity, linearity and distribution adjusted to normal (K-S test $p = 0.20$) were confirmed.

Comparing the models of HC and MDD patients, we can observe that the predictive variables: years of education, job status and physical activity, explain the 48% of variance of GCDI in MDD patients sample ($F=20.95$; $p<0.001$) compared to 24.9% in HC sample.

In the subsequent study of the interaction of the variables with the condition of being a MDD patients or HC subject, no interaction effects were observed with the variable of years of education ($F= 0.39$, $p= 0.89$) and physical activity ($F= 0.10$, $p= 0.99$).

Although, significant differences were observed between subjects who did little exercise and those who did it in a moderate way ($\bar{X}_i - \bar{X}= 0.42$, $p=0.04$) and in an intense way ($\bar{X}- \bar{X}= 0.48$, $p=0.03$) in the total sample (see Figure 2.).

On the other hand, an interaction of the condition with the job status was observed ($F=6.96$, $p=0.009$), there was a greater difference in means between active and nonactive patients than between active and inactive control subjects (see Figure 3.). In the sample of patients, an interaction between the age of illness onset and job status, was observed ($F=8.10$, $P= 0.008$). No interactions were observed in MDD patients between the age of onset of the disease and years of education ($F= 2.15$, $p= 0.18$) and physical activity ($F= 1.88$, $p=0.08$).

4. Discussion

Our results confirm that MDD patients exhibit global cognitive impairment and domain specific deficits, particularly on memory, latency of psychomotor speed and executive functions having problems in tasks of response to the item and information retrieval. Global cognitive deficits in MDD patients are associated with less years of education, lower physical activity, non-active job status and midlife and late onset of the disease. These variables have been widely related to each other (Vancamfort et al., 2015) and to cognitive performance (McDermott and Ebmeier, 2009; Snyder, 2013) but to date such a relationship had not been explored in MDD patients.

The age of illness onset was the only clinical variable associated with the GCDI in our study. This variable has been related in previous studies with a greater deterioration (Eraydin et al., 2018) and the changes in the cerebral anatomy (prefrontal cortex, hippocampus or the subcortical white matter) suffered by MDD patients (Kanner, 2004; Jannssen et al., 2007).

Traditionally, in the study of the illness onset have been taken into account the standards standard ranges of disease onset: <40 years (early onset), 40 to 60 years (midlife- onset) and >60 years (late-onset). These age ranges have been differentially

current diagnosis of Major Depressive Disorder

related to deficits in specific cognitive domains (Eraydin et al., 2018), observing in our study that the association between the global cognitive impairment increases progressively with the age of the disease.

The severity of the episode, a clinical factor that has been related in other studies with cognitive impairment (Yoshida et al., 2012), although inconsistently (McDermott and Ebmeier, 2009), has not been observed to be a clinical factor of global cognitive deterioration. This lack of agreement in the studies can be derived from the cognitive assessment tools and the methodology used to carry out the analyses. It is important that studies addressing the cognitive functioning in MDD are carried out in a more homogeneous way.

Years of schooling were also a significant factor associated with cognitive dysfunction of MDD patients and HC. In our sample, there was a significant difference between MDD patients and HC years of schooling, patients reported fewer years of regulated education than HC. Although the variables had not interaction, actions designed to increase the cognitive reserve of patients such as increasing reading hours, making logical games or learning new skills, should be useful. There is evidence that this construct may become a protective factor against potential cognitive dysfunction or may even compensate for cognitive competence when damage has already occurred (Venezia et al., 2018). In agreement with our results, Avila et al., found that elderly depressed patients with lower levels of education showed a worse performance in neuropsychological testing. By the contrary, other authors failed to obtain a relationship between cognitive performance and educational level in a sample of elderly patients (Bhalla et al, 2005), subjects with subclinical depressive symptoms

(McLaren et al, 2015) or in patients with MDD when the interaction with estimated intelligence was taking into account (Venezia et al, 2018).

People with a lower level of education are more vulnerable to factors such as unemployment, job insecurity or social status, which are factors associated to depression (Lorant et al., 2007; Bert et al., 2018). In our study, we have detected the risk of depressive patients to be in a non-active work situation due to unemployment or disability, which interacts significantly with their pathology, affecting both conditions in their cognitive deficit, which over time can be more difficult for this population group to reincorporate to an active labor status.

Regarding life-style factors, tobacco use has not been associated with GCDI, such as the study of Morisano et al. 2013, we haven't found a relationship between cognitive impairment and smoking status, but more research is needed. The new research should include larger sample sizes and longitudinal designs and to know more about this potential association evaluating the effects of smoking abstinence and reinstatement on cognition may be relevant.

On the other hand, physical activity was another significant factor related to cognitive impairment, with a higher probability of cognitive impairment for those who are more sedentary. In our sample we found significant differences between HC and patients in terms of physical activity; most MDD patients had little or no physical activity. It is known that there is a two-way relationship between physical activity and depression, people with depressive symptoms are more sedentary, due to lack of energy, apathy, physical comorbidity, and lower self-efficacy feelings (Vancampfort et al., 2015). On the other hand, a greater sedentary lifestyle may lead to depressive symptoms.

Individuals that exercise regularly show nearly 45% lower odds of having depressive symptoms than sedentary individuals, suggesting that low doses of activity can prevent depressive symptoms in healthy adults (Teychenne et al., 2008; Pareja-Galeano et al., 2016) and improve the course of MDD (Babyak et al., 2000). Physical activity has protective effects on the brain (Ferris et al., 2007; Gujral et al., 2017), exercise induce both acute and chronic effects affecting several putative systems implicated in depression such as neuroendocrine system, with the decrease in copeptin (Schuch et al, 2016) and cortisol (Beserra et al, 2018) neurogenesis, oxidative stress, with a marked reduction for thiobarbituric acid reactive substances (TBARS) was observed in a study of severely depressed inpatients, auto-immunity and producing cortical structural changes (Schuch et al, 2016). These results must be interpreted with caution since most biomarkers have been evaluated in only one or two studies and more studies are necessary.

Neural correlates of cognitive remediation have been poorly investigated. The few available studies have shown changes in cerebral activation after cognitive remediation: an increased activation in lateral and medial prefrontal, superior temporal, and lateral parietal regions (Meusel et al, 2013) and a trend toward normalization of intrinsic neural activity, with an increase in the right inferior frontal gyrus (a region that has been related to cognitive processing) (Schneider et al, 2019) and a modulation of left inferior parietal lobe and caudate nucleus activity, areas of interest in MDD.

Although it has not been demonstrated in a consistent way, in the meta-analysis of Sun et al., 2018, improvements were observed only in combination therapy with cognitive

therapy, MDD patients may benefit from practicing physical activity, because in addition to decreasing depressive symptoms, physical activity has been shown to be related to the improvement of cognitive performance, especially in older women (Hu et al., 2019). After a six-month physical activity intervention, Stillman et al. (2018) found improvements in memory and executive functioning in MDD patients.

It is important to understand the interrelation of variables and the importance of the study methodology of the studied variables and the population subject of study, when observing discrepancies in the works according to the age of the studied population, baseline cognitive alterations and clinical variables related to depression.

Finally, we have not found a relationship between cognitive deficit and metabolic syndrome. Unlike in other studies, we did not have to observe in our sample that people with metabolic syndrome have worse overall cognitive performance. This fact, may be due to the screening carried out in the exclusion criteria, in which subjects with certain comorbid diseases that may be related to the metabolic syndrome were eliminated. Therefore, we observed in our sample a small population suffering from metabolic syndrome.

Our study emphasizes the importance of taking cognitive deficits into account in depression. A high percentage of patients have domain-specific cognitive impairments and overall cognitive deficits, which affect their daily lives (Baune et al. 2010) and worsens their disease (Rock et al., 2014). We also highlight the importance of considering the different variables that can be associated with cognitive impairment in depression in order to be able to target these variables to prevent cognitive deficits.

In addition to being able to define these factors, it may open up the possibility of subsequently taking these variables into account in the treatment of cognitive symptoms of patients with depression for the study of pharmacological interventions that have not yet demonstrated their efficacy in the treatment of these symptoms (Devanand et al., 2018). In these studies it would be important to define samples that are as homogeneous as possible within the group and as heterogeneous as possible between the groups, differentiating patients with major depressive disorder and cognitive deficits from patients with mild cognitive impairment or patients with dementia.

Therefore, there is an important need to incorporate person-centred assessments and treatment protocols for the implementation of cost-effective and feasible augmentation strategies for the treatment of MDD. To this end, it is necessary to take into account the age at illness onset and to provide the patient with tools for an improvement in his cognitive reserve and to foster healthy lifestyles to prevent from cognitive deterioration. The accessible and well accepted behaviour such as walking 3-4 times weekly for 30-40 minutes during at least 9 weeks has positive effects on depressive symptoms, (Martinsen, 1995; Robertson et al., 2012, Kelly et al 2018) information which should be reported to all MDD patients in consultation, taking into account practical recommendations (Belvederi Murri et al., 2019) .

Our results highlight that physical activity can be a protective factor of cognitive functioning. It is relevant to investigate the preventive capacity of physical activity in the appearance of depression in vulnerable groups. Early and effective treatment in MDD is also important to prevent cognitive deterioration.

Limitations of the study include the relatively small sample size. Although we have used a small sample in the analysis, it is a heterogeneous population of MDD patients and HC. MDD patients have heterogeneous clinical scores, they have different grade of severity and duration of illness and treatment resistant depression scores.

Homogeneous scores have been obtained in the patient's and HC's BMI, therefore, we don't have representative scores for this variable. The cross-sectional design of the study is another of the limitations. More information on disturbances in cognitive performance and their possible relationships of dependency and interdependence with clinical and lifestyle factors can be obtained from longitudinal studies. Regarding data collection in the variables of healthy lifestyle, tobacco use and physical activity measures are self-reported. Self-reported measures may include biases in the study and in subsequent studies it would be advisable to objectively record both variables. On the one hand, taking records of the type of activity carried out by the subjects and time using heart rate monitors or accelerometers and, on the other hand, regarding tobacco use taking records of urinary cotinine concentrations and determination of exhaled carbon monoxide (eCO) level. Also, nicotine consumption was only collected through the number of cigarettes consumed per day. The use of other devices such as nicotine patches would be advisable in future studies. To perform the cognitive assessment, we used a validated tool. We explored different cognitive domains, but we did not perform cognitive auditory tasks or evaluated everyday cognitive skills.

5. Conclusions

Cognitive deficits were evident in patients with MDD compared with HC.

In our sample the GCDI of MDD patients can be explained by years of schooling, job status, physical activity and the age at illness onset. Although the model was also significant for HC, the explain variance was lower and the only significant variable was years of schooling. Therefore, these results support the importance of the implementation of interventions targeting the cognitive reserve and lifestyle habits of MDD patients, in addition to the conventional therapeutic approach focused on symptoms control.

Contributors

Pilar Lopez-García, Pilar Alvarez and Virginia Soria designed the study and wrote the protocol. Yolanda Sánchez-Carro and Maria J. Portella performed the statistical analyses. Yolanda Sánchez-Carro managed the literature searches and wrote the first draft of the manuscript, which was supervised by Pilar Lopez-García and Maria J. Portella. Itziar Leal-Leturia, Pilar Alvarez, Alba Toll, Virginia Soria, Neus Salvat-Pujol, Mikel Etxandi, Aida de Arriba-Arnau and Mikel Urretavizcaya participated in the recruitment. Yolanda Sánchez Carro, Pilar Alvarez, Esther Pousa, Alba Toll, Virginia Soria and Mikel Urretavizcaya administered the psychiatric interviews, psychometric scales and cognitive tests. All authors contributed to the interpretation and discussion of the results and have approved the final manuscript.

6.

7. acknowledgments

This study was supported in part by grants from the Carlos III Health Institute through the Ministry of Science, Innovation and Universities (PI15/00662, PI15/0039, PI15/00204), co-funded by the European Regional Development Fund (ERDF) “A way to

build Europe”, CIBERSAM, and the Catalan Agency for the Management of University and Research Grants (AGAUR 2017 SGR 1247). We also thank CERCA Programme/ Generalitat de Catalunya for institutional support. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. MJP is funded with a Miguel Servet II contract (CP16/00020) from the Carlos III Health Institute through the Spanish Ministry of Health. YSC work is supported by the FPI predoctoral grant (FPI 2016/17) from Universidad Autonoma de Madrid.

8. Conflict of Interest

The authors report no biomedical financial interests or potential conflicts of interest regarding this work.

9. References

1. World Health Organization. (2017). *Depression and other common mental disorders: global health estimates* (No. WHO/MSD/MER/2017.2).
2. Culpepper L, Muskin PR & Stahl SM (2015). Major depressive disorder: understanding the significance of residual symptoms and balancing efficacy with tolerability. *Am J Med*, 128 (9), S1-S15.
<https://doi.org/10.1016/j.amjmed.2015.07.001>.
3. Kennedy S (2002). Full remission: a return to normal functioning. *J Psychiatry Neurosci*, 27(4):233–4.
4. Lam RW, Kennedy SH, McIntyre RS, & Khullar A. (2014). Cognitive dysfunction in major depressive disorder: effects on psychosocial functioning and implications for treatment. *Can J Psychiatry*, 59(12), 649-654.

<https://doi.org/10.1177/070674371405901206>.

5. Rock PL, Roiser JP, Riedel WJ & Blackwell AD (2014). Cognitive impairment in depression: a systematic review and meta-analysis. *Psychol Med*, 44:2029–2040. <https://doi.org/doi: 10.1017/S0033291713002535>.
6. Baune BT & Air T (2016). Clinical, functional, and biological correlates of cognitive dimensions in major depressive disorder – rationale, design, and characteristics of the cognitive function and mood study (CoFaM-Study). *Front Psychiatry*; 7:150. <https://doi.org/ 10.3389/fpsy.2016.00150>.
7. Paelecke-Habermann Y, Pohl J & Leplo B (2005). Attention and executive functions in remitted major depression patients. *J affect disord*, 89(1-3), 125-135. <https://doi.org/10.1016/j.jad.2005.09.006>
8. Zheng H, Jia F, Guo G, Quan D, Li G, Wu H, Zhang B, Fan C, He X, & Huang, H. (2015). Abnormal anterior cingulate N-acetylaspartate and executive functioning in treatment-resistant depression after rTMS therapy. *Int J Neuropsychopharmacol*, 18(11). <https://doi.org/ 10.1093/ijnp/pyv059>.
9. Eraydin IE, Mueller C, Corbett A, Ballard C, Brooker H, Wesnes K, Aarsland D, & Huntley J. (2019). Investigating the relationship between age of onset of depressive disorder and cognitive function. *Int J Geriatr Psychiatry*, 34(1), 38-46.

<https://doi.org/10.1002/gps.4979>.
10. Yoshida T, Ishikawa M, Niitsu T, Nakazato M, Watanabe H, Shiraishi T, Shijna A, Hashimoto T, Kanahara N, Hasegawa T, Enohara M, Kimura A, Iyo M & Hashimoto K (2013). * Decreased serum levels of mature brain-derived

neurotrophic factor (BDNF), but not its precursor proBDNF, in patients with major depressive disorder [published correction appears in PLoS One.

2013;8(2)]. *PLoS One*. 2012;7(8):e42676.

<https://doi.org/10.1371/journal.pone.0042676>

11. Kanner AM (2004). Structural MRI changes of the brain in depression. *Clin EEG Neurosci*, 35(1), 46-52. <https://doi.org/10.1177/155005940403500111>.
12. Janssen J, Hulshoff Pol HE, de Leeuw FE, Schnack HG, Lampe IK, Kok RM, Kahn RS & Heeren TJ (2007). Hippocampal volume and subcortical white matter lesions in late life depression: comparison of early and late onset depression. *J Neurol Neurosurg Psychiatry*;78(6):638–40. <https://doi.org/10.1016/j.biopsych.2004.09.011>
13. Salvat-Pujol N, Labad J, Urretavizcaya M, de Arriba-Arnau A, Segalàs C, Real E, Ferrer A, Crespo JM, Jiménez-Murcia S, Soriano-Mas C, Menchón JM, Soria V (2017). Hypothalamic pituitary-adrenal axis activity and cognition in major depression: The role of remission status. *Psychoneuroendocrinology*, 76:38-48. <https://doi.org/10.1016/j.psyneuen.2016.11.007>.
14. Chen Y, Wang C, Zhu X, Tan Y, & Zhong Y (2015). Aberrant connectivity within the default mode network in first-episode, treatment major depressive disorder. *J affect disord*, 183, 49-56. <https://doi.org/10.1016/j.jad.2015.04.052>
15. Douglas KM & Porter RJ (2009). Longitudinal assessment of neuropsychological function in major depression. *Aus N Z J Psychiatry*, 43(12), 1105-1117. <https://doi.org/10.3109/00048670903279887>.

16. Davis M, DellaGioia N, Matuskey D, Harel B, Maruff P, Pietrzak RH & Esterlis, I (2017). Preliminary evidence concerning the pattern and magnitude of cognitive dysfunction in major depressive disorder using cogstate measures. *J Affect Disord*, 218, 82. <https://doi.org/10.1016/j.jad.2017.04.064>.
17. Depp CA, Bowie CR, Mausbach BT, Wolyniec P, Thornquist MH, Luke JR, McGrath JA, Pulver AE, Patterson TL, Harvey PD (2015). Current smoking is associated with worse cognitive and adaptive functioning in serious mental illness. *Acta Psychiatr Scand*, 131(5):333-41. <https://doi.org/10.1111/acps.12380>.
18. Aas M, Djurovic S, Ueland T, Mørch RH, Fjæra Laskemoen J, Reponen EJ, Cattaneo A, Eiel Steen N, Agartz I, Melle I, Andreassen OA (2019). *The relationship between physical activity, clinical and cognitive characteristics and BDNF mRNA levels in patients with severe mental disorders*. *World J Biol Psychiatry*, 20(7):567-576. <https://doi.org/10.1080/15622975.2018.1557345>.
19. Breslau, N, Kilbey, MM, & Andreski, P (1993). Nicotine dependence and major depression: new evidence from a prospective investigation. *Archives of general psychiatry*, 50(1), 31-35. <https://doi.org/10.1001/archpsyc.1993.01820130033006>
20. Pareja-Galeano H, Mayero S, Perales M, Garatachea N, Santos-Lozano A, Fiuza-Luces C, Emanuele E, Gálvez BG, Sanchis-Gomar F & Lucia A (2016). Biological rationale for regular physical exercise as an effective intervention for the prevention and treatment of depressive disorders. *Curr Pharm Des*, 22(24), 3764-3775. <https://doi.org/10.2174/1381612822666160322144537>.

21. Kramer AF, & Erickson KI (2007). Capitalizing on cortical plasticity: Influence of physical activity on cognition and brain function. *Trends Cogn Sci*, 11(8), 342–348. <https://doi:10.1016/j.tics.2007.06.009>.
22. Cassilhas RC, Tufik S, & de Mello MT (2016). Physical exercise, neuroplasticity, spatial learning and memory. *Cell Mol Life Sci*, 73(5), 975-983. <https://doi.org/10.1007/s00018-015-2102-0>.
23. Karssemeijer EE, Aaronson JJ, Bossers WW, Smits TT, & Kessels RR (2017). Positive effects of combined cognitive and physical exercise training on cognitive function in older adults with mild cognitive impairment or dementia: A meta-analysis. *Ageing Res Rev*, 40, 75-83. <https://doi.org/10.1016/j.arr.2017.09.003>.
24. Schuch FB, Deslandes AC, Stubbs B, Gosmann NP, Silva CTB & Fleck MPA (2015) Neurobiological effects of exercise on Major Depressive Disorder: A systematic review. *Neurosci Biobehav Rev*; 61. 1-11. <http://dx.doi.org/10.1016/j.neubiorev.2015.11.012>
25. Lambourne K, Audiffren M, & Tomporowski PD (2010). Effects of acute exercise on sensory and executive processing tasks. *Med Sci Sports Exerc*, 42, 1396–1402. <https://doi.org/10.1249/MSS.0b013e3181cbee11>.
26. McMorris T, Davranche K, Jones G, Hall B, & Minter C. (2009). Acute incremental exercise, performance of a central executive task, and sympathoadrenal system and hypothalamic–pituitary–adrenal axis activity. *Int J Psychophysiol*; 73 (3): 334-340. <https://doi.org/10.1016/j.ijpsycho.2009.05.004>.

27. Ferris LT, Williams JS, & Shen C. (2007). The effect of acute exercise on serum brain-derived neurotrophic factor levels and cognitive function. *Med Sci Sports Exerc*, 39, 728–734. <https://doi.org/10.1249/mss.0b013e31802f04c7>.
28. Meusel LAC, Hall GBC, Fougere P, McKinnon MC, MacQueen GM. Neural Correlates of Cognitive Remediation in Patients With Mood Disorders. *Psychiatry Res*. 2013 30;214(2):142-52. <https://doi.org/10.1016/j.psychresns.2013.06.007>.
29. Schneider I, Schmitgen MM, Bach C, Listunova L, Kienzle J, Sambataro F, Wolf RC (2019). Cognitive remediation therapy modulates intrinsic neural activity in patients with major depression. *Psychological Medicine*, 2019, 1–11. <https://doi.org/10.1017/s003329171900240x>.
30. Gujral S, Aizenstein H, Reynolds III CF, Butters MA, & Erickson KI (2017). Exercise effects on depression: possible neural mechanisms. *Gen Hosp Psychiatry*, 49, 2-10. <https://doi.org/10.1016/j.genhosppsy.2017.04.012>.
31. Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ, Webb A, Jerome GJ, Marquez DX & Elavsky S (2004). Cardiovascular fitness, cortical plasticity and aging. *Proc Natl Acad Sci USA*; 101, 3316–3321. <https://doi.org/10.1073/pnas.0400266101>
32. Erickson KI, Voss MW, Prakash RS Basak C, Szabo A, Chaddock L, Kim JS, Heo S, Alves H, White SM, Wojcicki TR, Mailey E, Vieira VJ, Martin SA, Pence BD, Woods JA, McAuley E & Kramer AF (2011) .Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci USA*; 108(7):3017–3022. <https://doi.org/10.1073/pnas.1015950108>.

33. Kubesch S, Breschneider V, Freudenmann R, Weidenhammer N, Lehmann M, Spitzer M & Grön G (2003). Aerobic endurance exercise improves executive functions in depressed patients. *J Clin Psychiatry*, 64, 1005 – 1012.
[https://doi.org/ 10.4088/jcp.v64n0905](https://doi.org/10.4088/jcp.v64n0905)
34. Colcombe S, & Kramer AF (2003) Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci*. 14, 125–130.
<https://doi.org/10.1111/1467-9280.t01-1-01430>
35. Rathore A, & Lom B (2017). The effects of chronic and acute physical activity on working memory performance in healthy participants: a systematic review with meta-analysis of randomized controlled trials. *Syst Rev*, 6(1), 124.
<https://doi.org/10.1186/s13643-017-0514-7>.
36. Heishman SJ, Kleykamp BA & Singleton EG (2010). Meta-analysis of the acute effects of nicotine and smoking on human performance. *Psychopharmacology (Berl)*; 210 (4): 453 – 469. <https://doi.org/10.1007/s00213-010-1848-1>.
37. Durazzo TC, Meyerhoff DJ & Nixon SJ (2010). Chronic cigarette smoking: Implications for neurocognition and brain neurobiology. *Int J Environ Res Public Health*; 7:3760–3791. <https://doi.org/10.3390/ijerph7103760>.
38. Wagner M, Schulze-Rauschenbach S, Petrovsky Brinkmeyer J, von der Goltz C, Gründer G, Spreckelmeyer KN, Wienker T, Diaz-Lacava A, Mobascher A, Dahmen N, Clepce M, Thuerauf N, Kiefer F, de Millas JW, Gallinat J, Winterer G (2012). Neurocognitive impairments in non-deprived smokers-results from a population-based multi-center study on smoking-related behaviour. *Addict Biol*; 18 (4):752, 761. <https://doi.org/10.1111/j.1369-1600.2011.00429.x>.

39. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D & Bor DH (2000). Smoking and mental illness: A population-based prevalence study. *JAMA*. 284 (20) :2606–2610. <https://doi.org/10.1001/jama.284.20.2606>.
40. Prochaska JJ (2011). Smoking and mental illness—breaking the link. *N Engl J Med*, 365(3), 196-198. <https://doi.org/10.1056/NEJMp1105248>.
41. Weinberger AH, Kashan RS, Shpigel DM, Esan H, Taha F, Lee CJ, Funk AP & Goodwin RD (2017). Depression and cigarette smoking behaviour: a critical review of population-based studies. *Am J Drug Alcohol Abuse*, 43(4), 416-431. <https://doi.org/10.3109/00952990.2016.1171327>.
42. Morisano D, Wing VC, Sacco KA, Arenovich T & George TP (2013). Effects of tobacco smoking on neuropsychological function in schizophrenia in comparison to other psychiatric disorders and non-psychiatric controls. *American J Addict* 22(1), 46-53. <https://doi.org/10.1111/j.1521-0391.2013.00313.x>.
43. Cleeman JL, Grundy SM, Becker D & Clark L (2001). Expert on detection evaluation and treatment of high blood cholesterol in adults. Executive summary of the third report of the National Cholesterol Education Program (NCEP) Adult Treatment Panel (ATP III) *Jama*, 285 (19), 2486-2497.
44. Mitchell AJ, Vancampfort D, Sweers K, van Winkel R, Yu W & De Hert M (2011). Prevalence of metabolic syndrome and metabolic abnormalities in schizophrenia and related disorders—a systematic review and meta-analysis. *Schizophr bull*, 39(2), 306-318. <https://doi.org/10.1093/schbul/sbr148>

45. Vancampfort D, Vansteelandt K, Correll CU, Mitchell AJ, De Herdt A, Sienaert P, Probst M & De Hert M. (2013). Metabolic syndrome and metabolic abnormalities in bipolar disorder: a meta-analysis of prevalence rates and moderators. *Am J Psychiatry*, 170(3), 265-274.
<https://doi.org/10.1176/appi.ajp.2012.12050620>.
46. Vancampfort D, Correll CU, Wampers M, Sienaert P, Mitchell AJ, De Herdt A, Probst M, Scheewe TW, & De Hert M. (2014). Metabolic syndrome and metabolic abnormalities in patients with major depressive disorder: a meta-analysis of prevalences and moderating variables. *Psychol Med*, 44(10), 2017-2028. <https://doi.org/10.1017/S0033291713002778>.
47. Vogelzangs N, Beekman AT, van Reedt Dortland AK, Schoevers RA, Giltay EJ, de Jonge P & Penninx BW (2014). Inflammatory and metabolic dysregulation and the 2-year course of depressive disorders in antidepressant users. *Neuropsychopharmacology*, 39(7):1624–1634.
<https://doi.org/10.1038/npp.2014.9>
48. Marijnissen RM, Vogelzangs N, Mulder ME, van den Brink RH, Comijs HC & Oude Voshaar RC (2017). Metabolic dysregulation and late-life depression: a prospective study. *Psychol Med*; 47 (6):1041-1052.
<https://doi.org/10.1017/S0033291716003196>.
49. Pan A, Keum N, Okereke OI, Sun Q, Kivimaki M, Rubin RR & Hu FB (2012). Bidirectional association between depression and metabolic syndrome: a systematic review and meta-analysis of epidemiological studies. *Diabetes Care*;35(5):1171–1180. <https://doi.org/10.2337/dc11-2055>

50. Livingston JM, McDonald MW, Gagnon T, Jeffers MS, Gomez-Smith M, Antonescu S, Cron GO, Boisvert C, Lacoste B, Corbett D (2020). Influence of metabolic syndrome on cerebral perfusion and cognition. *Neurobiol Dis*; 137:104756. <https://doi.org/10.1016/j.nbd.2020.104756>.
51. Danna SM, Graham E, Burns RJ, Deschenes SS & Schmitz N (2016). Association between depressive symptoms and cognitive function in persons with diabetes mellitus: a systematic review. *PLoS One*; 11: e0160809. <https://doi.org/10.1371/journal.pone.0160809>.
52. Sullivan MD, Katon WJ, Lovato LC, Miller ME, Murray AM, Horowitz KR, Nick Bryan R, Gerstein HC, Marcovina S, M.D., Akpunonu BE, Johnson J, Yale JF, Williamson J, & Launer LJ (2013). Association of depression with accelerated cognitive decline among patients with type 2 diabetes in the ACCORD-MIND trial. *JAMA Psychiatry*; 70: 1041–7. <https://doi.org/10.1001/jamapsychiatry.2013.1965>.
53. Allan JL, McMinn D, Daly M (2016). A Bidirectional Relationship between Executive Function and Health Behavior: Evidence, Implications, and Future Directions. *Front Neurosci*; 10: 386. <https://doi.org/10.3389/fnins.2016.00386>
54. Daly M, McMinn M, Allan JL (2014). A bidirectional relationship between physical activity and executive function in older adults. *Front Hum Neurosci*; 8: 1044. <https://doi.org/10.3389/fnhum.2014.01044>.
55. American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text rev.). Washington, DC: American Psychiatric Association.

56. Fekadu A, Wooderson SC, Markopoulou K & Cleare AJ (2009). The Maudsley Staging Method for treatment-resistant depression: prediction of longer-term outcome and persistence of symptoms. *J Clin Psychiatry*, 70(7), 952-957. <https://doi.org/10.4088/JCP.08m04728>.
57. Fekadu, A., Donocik, J.G. & Cleare, A.J (2018) Standardisation framework for the Maudsley staging method for treatment resistance in depression. *BMC Psychiatry* 18, 100. <https://doi.org/10.1186/s12888-018-1679-x>.
58. Levaux MN, Potvin S, Sepehry AA, Sablier J, Mendrek A, & Stip E (2007). Computerized assessment of cognition in schizophrenia promises and pitfalls of CANTAB. *Eur Psychiatry*, 22(2), 104-115. <https://doi.org/10.1016/j.eurpsy.2006.11.004>.
59. Kim HS, An YM, Kwon JS, & Shin MS (2014). A preliminary validity study of the cambridge neuropsychological test automated battery for the assessment of executive function in schizophrenia and bipolar disorder. *Psychiatry Investig*, 11(4), 394. <https://doi.org/10.4306/pi.2014.11.4.394>.
60. Sweeney JA, Kmiec JA & Kupfer DJ (2000). Neuropsychologic impairments in bipolar and unipolar mood disorders on the CANTAB Neurocognitive Battery. *Biol Psychiatry*, 48: 674-685. [https://doi.org/10.1016/s0006-3223\(00\)00910-0](https://doi.org/10.1016/s0006-3223(00)00910-0).
61. Vancampfort D, Stubbs B, Sienaert P, Wyckaert S, De Hert M, Rosenbaum S & Probst M (2015). What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. *Psychiatr Danub*; 27(3):210-24.

62. McDermott LM, & Ebmeier KP (2009). A meta-analysis of depression severity and cognitive function. *J Affect Disord* 119(1-3), 1-8.
<https://doi.org/10.1016/j.jad.2009.04.022>.
56. Snyder HR (2013). Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: a meta-analysis and review. *Psychol Bull*, 139(1), 81.
<https://doi.org/10.1037/a0028727>.
57. Janssen J, Hilleke E, Pol H, Lampe IK, Schnack HG, de Leeuw FK, Kahn RS & Heeren TJ (2004). Hippocampal changes and white matter lesions in early-onset depression. *Biol Psychiatry*. 2004;56(11):825-831.
<https://doi.org/10.1016/j.biopsych.2004.09.011>.
58. McLaren ME, Szymkowicz SM, Kirton JW & Dotson VM (2015). Impact of education on memory deficits in subclinical depression. *Archives of Clinical Neuropsychology*. 2015 30: 387-393. <https://doi.org/10.1093/arclin/acv038>.
59. Venezia RG, Gorlyn M, Burke AK, Oquendo MA, Mann JJ & Keilp JG (2018). The impact of cognitive reserve on neurocognitive performance in Major Depressive Disorder. *Psychiatry res*, 270, 211-218.
<https://doi.org/10.1016/j.psychres.2018.09.031>
60. Avila R, Moscoso MA, Ribeiz S, Arrais J, Jaluul O & Bottino CM (2009). Influence of education and depressive symptoms on cognitive function in the elderly. *Int Psychogeriatr*;21(3):560-7. <https://doi.org/10.1017/S1041610209008928>.
61. Bhalla RK, Butters MA, Zmuda MD, Seligman K, Mulsant BH, Pollock BG, & Reynolds CF (2005). Does education moderate neuropsychological impairment

- in late-life depression? *International Journal of Geriatric Psychiatry*, 20(5), 413–417. <https://doi.org/10.1002/gps.1296>.
62. Lorant V, Croux C, Weich S, Delière D, Mackenbach J & Ansseau M (2007). Depression and socio-economic risk factors: 7-year longitudinal population study. *Br j psychiatry*, 190(4), 293-298. <https://doi.org/10.1192/bjp.bp.105.020040>
63. Bert F, Gualano MR, Thomas R, Vergnano G, Voglino G & Siliquini R (2018). Exploring the possible health consequences of job insecurity: a pilot study among young workers. *Gac sanit*. <https://doi.org/10.1016/j.gaceta.2018.08.011>
64. Teychenne M, Ball K & Salmon J (2008). Physical activity and likelihood of depression in adults: a review. *Prev Med*, 46(5), 397-411. <https://doi.org/10.1016/j.ypmed.2008.01.009>.
65. Babyak M, Blumenthal JA, Herman S, Khatrri P, Doraiswamy M, Moore K, Craighead WE, Baldewicz TT & Krishnan KR (2000). Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom med*, 62(5), 633-638. <https://doi.org/10.1097/00006842-200009000-00006>.
66. Beserra AHN, Kameda P, Deslandes AC, Schuch FB, Laks J, Moraes HS (2018). Can physical exercise modulate cortisol level in subjects with depression? A systematic review and meta-analysis. *Trends Psychiatry Psychother*.;40(4):360-8. <https://doi.org/10.1590/2237-6089-2017-0155>.
67. Sun M, Lanctot K, Herrmann N & Gallagher D (2018). Exercise for Cognitive Symptoms in Depression: A Systematic Review of Interventional Studies. *Can J Psychiatry* 63, 115-128.

<https://doi.org/10.1177/0706743717738493>.

68. Hu, L., Smith, L., Imm, K.R., Jackson, S.E., and Yang, L. (2019). Physical activity modifies the association between depression and cognitive function in older adults. *J Affect Disord* 246, 800-805. <https://doi.org/10.1016/j.jad.2019.01.008>
69. 69. Stillman CM, & Erickson KI (2018). Physical activity as a model for health neuroscience. *Ann N Y Acad Sci*, 1428(1), 103-111.
<https://doi.org/10.1111/nyas.13669>
70. Baune BT, Miller R, McAfoose J, Johnson M, Quirk F, & Mitchell D (2010). The role of cognitive impairment in general functioning in major depression. *Psychiatry res*, 176(2-3), 183-189.
<https://doi.org/10.1016/j.psychres.2008.12.001>.
71. Devanand DP, Pelton GH, D'Antonio K, Ciarleglio A, Scodes J, Andrews H, Lunsford J, Beyer JL, Beyer JR, Sneed J, Ciovacco M & Doraiswamy PM (2018). Donepezil Treatment in Patients With Depression and Cognitive Impairment on Stable Antidepressant Treatment: A Randomized Controlled Trial. *Am J Geriatr Psychiatry*;26(10):1050-1060. <https://doi.org/10.1016/j.jagp.2018.05.008>
72. Martinsen EW (1995). The effects of exercise on mental health in clinical populations. *European perspectives on exercise and sport psychology.*, 71-84.
73. Robertson R, Robertson A, Jepson R & Maxwell M (2012). Walking for depression or depressive symptoms: a systematic review and meta-analysis. *Mental health and physical activity*, 5(1), 6675.
<https://doi.org/10.1016/j.mhpa.2012.03.002>

74. Kelly P, Williamson C, Niven AG, Hunter R, Mutrie N & Richards J (2018).

Walking on sunshine: scoping review of the evidence for walking and mental health. *Br J sports med*, 52(12), 800-806. <https://doi.org/10.1136/bjsports-2017-098827>.

75. Belvederi Murri M, Ekkekakis P, Magagnoli M, Zampogna D, Cattedra S,

Capobianco L, Serafini G, Calcagno P & Amore, M. (2019). Physical exercise in major depression: reducing the mortality gap while improving clinical outcomes. *Frontiers in psychiatry*, 9, 762. <https://doi.org/10.3389/fpsyt.2018.00762>

Tables and figures

Table 1. Demographics, clinical, metabolic and health-related variables of HC and MDD patients.

Variable	HC N = 68	MDD N = 74	t/ χ^2	p
Age	48.28 (10.25)	49.73 (10.26)	0.71	0.4
Gender (% of males)	29.4	29.7	0.25	0.6
Years of education	15.81 (5.70)	13.18 (4.59)	9.09	0.003
Premorbid IQ	115.32 (10.85)	109.65(11.80)	0.48	0.003
Marital Status (%)			2.01	0.1
Single	25	28.8		
Partnered / Married	60.3	54.8		
Separated/ Divorced	8.8	12.3		
Widowed	5.9	4.1		
Current work status (%)			49.12	<0.001
Active	89.7	28.4		
Non- active:	10.3	71.6		

current diagnosis of Major Depressive Disorder

Unemployed	8.8	18.9		
Disability	1.5	47.3		
Retired	0	5.4		
Metabolic Syndrome (%)	23.1	16.2	1.04	0.31
BMI	26.36 (4.64)	26.44 (4.55)	0.13	0.9
FBS	95.33 (13.21)	96.99 (17.38)	1.95	0.5
HDL-C	72.16(42.13)	68.50 (28.57)	1.97	0.5
Triglyceride level	115.14 (82.46)	99.63 (57.33)	4.68	0.2
Waist measure	88.27 (13.60)	90.71 (12.26)	1.26	0.3
SBP	127.70 (17.65)	123.86 (17.78)	1.64	0.2
DBP	78.63 (9.96)	76.61 (12.09)	1.15	0.2
Tobacco use (%)			2.92	0.2
Smokers	19.5	29.8		
Abstainers	29.8	37.8		
Non-smokers	50.7	32.4		
Physical activity (%)			6.60	0.03
Low	19.4	39.2		
Moderate	50.7	39.2		
High	29.9	21.6		
WHODAS	13.48 (2.11)	33.95 (10.11)	263.99	<0.001
S-Anxiety	9.60 (6.47)	37.37 (11.38)	307.81	<0.001
Variable	HC N = 68	MDD N = 74	t/ χ^2	p
MSM (%)				
Mild		12.2		
Moderate		56.8		
Severe		31.1		
HDRS- 17		20.01 (5.41)		
Age at illness onset		32.74 (13.72)		
Antidepressant medication (%)				
SSRI		32.4		
SNRIs		33.8		
TCAs		9.5		
TeCAs		17.6		

Footnote 1. Values represent mean and standard deviation (SD)

HC= Healthy controls; MDD= Major Depressive Disorder patients.

IQ= Intelligence Quotient, WHODAS= World Health Organization Disability Assessment

Schedule 2.0., S-Anxiety= State Anxiety Inventory, T- Anxiety= Trait Anxiety Inventory, BMI=

body mass index, FBS= Fasting Blood Sugar level, HDL-C= High-Density Lipoprotein Cholesterol,

current diagnosis of Major Depressive Disorder

SBP= Systolic Blood Pressure, DBP= Diastolic Blood Pressure, MSM =Maudsley Staging Method, HDRS-17= Hamilton Depression Rating Scale-17, SRIs= Serotonin Reuptake Inhibitors, SNRIs= Serotonin-Norepinephrine Reuptake Inhibitors, TCAs= Tricyclic Antidepressants, TeCAs= Tetracyclic antidepressants .

Journal Pre-proof

current diagnosis of Major Depressive Disorder

Table 2. Cognitive scores of HC and MDD patients.

Cognitive Variables	HC N = 68	MDD N = 74	F	P
<i>Attention</i>				
RVP total hits	20.71 (9.83) -0.22 (0.92)	16.40 (9.45) 0.21 (1.03)	6.89	0.010
<i>Memory</i>				
DMS correct trials	25.37 (3.83) -0.23 (0.91)	21.54 (5.32) 0.21 (1.04)	7.10	<0.001 0.009
<i>Executive function</i>				
SWM strategy	32.31 (4.93) -0.15 (0.97)	34.53 (5.30) 0.14(1.01)	3.13	0.07
OTS correct on first choice	14.09 (5.08) -0.26 (0.92)	11.61 (5.69) 0.24 (1.02)	9.13	0.006 0.003
<i>Latency</i>				
MTS latency	707.72 (167.89) -0.24 (0.90)	869.94 (327.75) 0.22(1.04)	8.04	<0.001 0.005
SWM time to last response	27387.50 (7460.68) -0.16 (0.96)	32445.05 (9655.10) 0.15(1.02)	3.43	0.001 0.06
<i>Impulsivity</i>				
RVP false alarm	2.07 (3.19) -0.20 (0.94)	2.77 (4.59) 0.18 (1.02)	5.19	0.024
OTS false alarm	-0.08 (0.38) -0.22 (0.97)	0.07 (0.75) 0.20 (0.99)	6.75	0.010
GCDI	-0.24 (0.90)	0.22 (1.04)	8.29	0.005

Footnote 2. Values represent mean and standard deviation (SD) of raw scores and adjusted z-scores.

P values of raw scores included were deemed significant following corrections for multiple testing.

Cognitive z-scores (RVP, DMS, SWM, OTS, MOT and Global Cognitive Dysfunction Index) have been adjusted by age, IQ and Gender to calculate the statistical value and its significance.

HC= Healthy controls; MDD= Major depressive Disorder patients. RVP= Rapid visual information process test, DMS= Delayed Matching to Sample, SWM= Spatial working memory, OTS= One touch Stockings of Cambridge, MOT= Motor Screening Task, GCDI = Global Cognitive Dysfunction Index.

current diagnosis of Major Depressive Disorder

Figure 1. Scatterplot of multivariate regression model.

Footnote 3. *The horizontal axis indicates the predictive value of all included variables in the regression model. The vertical axis indicates the measured value of GCDI. The dots indicate each patient. The red line corresponds to the regression line of the model.*

GCDI = Global Cognitive Dysfunction Index.

Figure 2. Global Cognition Dysfunction Scores by Physical Activity.

Footnote 4. *The horizontal axis indicates the physical activity levels of the sample: Low, moderate and high measures by IPAQ scale. The vertical axis indicates the measured value of GCDI. The columns indicate the means of each group and the line indicates the means of the total sample.*

GCDI = Global Cognitive Dysfunction Index.

current diagnosis of Major Depressive Disorder

Figure 3. Global Cognition Dysfunction Scores by job status in HC and MDD patients

Footnote 5. *The horizontal axis indicates the condition of the sample: healthy controls or Major Depressive Disorder patients. The vertical axis indicates the measured value of GCDI. The asterisk indicates the differences in scores between groups that were statistically significant. GCDI = Global Cognitive Dysfunction Index.*