

**Association of Prenatal Exposure to Persistent Organic
Pollutants with Obesity and Cardiometabolic Traits
in Early Childhood: The Rhea Mother–Child Cohort
(Crete, Greece)**

**Marina Vafeiadi, Vaggelis Georgiou, Georgia Chalkiadaki,
Panu Rantakokko, Hannu Kiviranta, Marianna Karachaliou,
Eleni Fthenou, Maria Venihaki, Katerina Sarri, Maria Vassilaki,
Soterios A. Kyrtopoulos, Emily Oken, Manolis Kogevinas,
and Leda Chatzi**

<http://dx.doi.org/10.1289/ehp.1409062>

Received: 11 August 2014

Accepted: 21 April 2015

Advance Publication: 24 April 2015

This article will be available in its final, 508-conformant form 2–4 months after Advance Publication. If you need assistance accessing this article before then, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Association of Prenatal Exposure to Persistent Organic Pollutants with Obesity and Cardiometabolic Traits in Early Childhood: The Rhea Mother–Child Cohort (Crete, Greece)

Marina Vafeiadi,¹ Vaggelis Georgiou,¹ Georgia Chalkiadaki,¹ Panu Rantakokko,² Hannu Kiviranta,² Marianna Karachaliou,¹ Eleni Fthenou,¹ Maria Venihaki³, Katerina Sarri,¹ Maria Vassilaki,¹ Soterios A. Kyrtopoulos,⁴ Emily Oken,⁵ Manolis Kogevinas,^{6,7,8,9} and Leda Chatzi¹

¹Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Greece;

²Department of Environmental Health, National Institute for Health and Welfare, Kuopio,

Finland; ³Lab of Clinical Chemistry-Biochemistry, Department of Laboratory Medicine, School

of Medicine, University of Crete, Heraklion, Crete, Greece; ⁴National Hellenic Research

Foundation, Institute of Biology, Medicinal Chemistry and Biotechnology, Athens, Greece;

⁵Obesity Prevention Program, Department of Population Medicine, Harvard Medical School and

Harvard Pilgrim Health Care Institute, Boston, Massachusetts, USA; ⁶Centre for Research in

Environmental Epidemiology (CREAL), Barcelona, Spain; ⁷Hospital del Mar Research Institute

(IMIM), Barcelona, Spain; ⁸CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona,

Spain; ⁹National School of Public Health, Athens, Greece

Address correspondence to Marina Vafeiadi, Department of Social Medicine, Faculty of

Medicine, University of Crete, P.O. Box 2208, Heraklion 71003, Crete, Greece. E-mail:

m.vafeiadi@med.uoc.gr

Running title: POPs, obesity and cardiometabolic risk

Acknowledgments: The authors would particularly like to thank all the cohort participants for their generous collaboration. The Rhea project was financially supported by European projects (EU FP6-2003-Food-3-NewGeneris, EU FP6. STREP Hiwate, EU FP7 ENV.2007.1.2.2.2. Project No 211250 Escape, EU FP7-2008-ENV-1.2.1.4 Envirogenomarkers, EU FP7-HEALTH-2009- single stage CHICOS, EU FP7 ENV.2008.1.2.1.6. Proposal No 226285 ENRIECO, EU-FP7- HEALTH-2012 Proposal No 308333 HELIX), MeDALL (FP7 European Union project, No. 264357), and the Greek Ministry of Health (Program of Prevention of obesity and neurodevelopmental disorders in preschool children, in Heraklion district, Crete, Greece: 2011-2014; “Rhea Plus”: Primary Prevention Program of Environmental Risk Factors for Reproductive Health, and Child Health: 2012-15). Dr. Oken was supported by the US National Institutes of Health (K24 HD069408).

Competing financial interests: The authors declare they have no competing financial interests.

Abstract

Background: Prenatal exposure to endocrine disrupting chemicals such as persistent organic pollutants (POPs) may increase risk of obesity later in life.

Aims: We examined the relation of *in utero* POPs exposure to offspring obesity and cardiometabolic risk factors at 4 years in the Rhea mother-child cohort in Crete, Greece (n=689).

Methods: We determined concentrations of polychlorinated biphenyls (PCBs), dichlorodiphenyldichloroethene (DDE), and hexachlorobenzene (HCB) in first trimester maternal serum. We measured child weight, height, waist circumference, skinfold thicknesses, blood pressure (BP), blood levels of lipids, C-reactive protein, and adipokines at 4 years of age. Childhood obesity was defined using age- and sex-specific cut points for BMI as recommended by the International Obesity Task Force.

Results: On multivariable regression analyses, a 10-fold increase in HCB was associated with a higher BMI z score (adj. $\beta=0.49$; 95% CI: 0.12, 0.86), obesity (RR=8.14; 95% CI: 1.85, 35.81), abdominal obesity (RR=3.49; 95% CI: 1.08, 11.28), greater sum of skinfold thickness ($\beta=7.71$ mm; 95% CI: 2.04, 13.39) and higher systolic BP ($\beta=4.34$ mmHg; 95% CI: 0.63, 8.05) at 4 years of age. Prenatal DDE exposure was associated with higher BMI z score ($\beta=0.27$; 95% CI: 0.04, 0.5), abdominal obesity (RR=3.76; 95% CI: 1.70, 8.30) and higher diastolic BP ($\beta=1.79$ mmHg; 95% CI: 0.13, 3.46). PCBs were not significantly associated with offspring obesity or cardiometabolic risk factors.

Conclusions: Prenatal exposure to DDE and HCB was associated with excess adiposity and higher blood pressure levels in early childhood.

Introduction

Persistent organic pollutants (POPs) are ubiquitous and persist in the environment, accumulate in high concentrations in fatty tissues, and are bio-magnified through the food-chain. POPs include synthetic chemicals that were widely used as pesticides [e.g., dichlorodiphenyltrichloroethane (DDT), dichlorodiphenyldichloroethene (DDE), hexachlorobenzene (HCB)] and in industrial processes [polychlorinated biphenyls (PCBs)], through most of the 20th century. Human exposure to POPs occurs primarily through diet, and maternal concentrations of POPs are transmitted to the developing fetus prenatally through the placenta and postnatally via breast milk (Needham et al. 2011). The use of these chemicals is presently banned (PCBs, HCB) or restricted (DDT) (Stockholm Convention on Persistent Organic Pollutants 2009). However, because of their persistence in the environment, the general population is still exposed to these substances at low doses (Porta et al. 2008) and adverse health outcomes related to background levels of exposure are still a concern (WHO 2010). Moreover, several of these compounds act as endocrine disrupting chemicals (EDCs) which can alter the normal function of endocrine systems in humans and wildlife (Diamanti-Kandarakis et al. 2009).

In utero exposure to EDCs has been hypothesized to increase risk of obesity in childhood and into adulthood (Baillie-Hamilton 2002; Diamanti-Kandarakis et al. 2009; La Merrill and Birnbaum 2011). According to the “environmental obesogen hypothesis”, early-life exposure to chemicals that could mimic or block the natural action of endogenous hormones may perturb the mechanisms involved in adipogenesis or energy storage and, thus, may increase an individual’s susceptibility to obesity (Grun and Blumberg 2006).

Several longitudinal birth cohort studies have examined the relationship of prenatal exposure to EDCs with child growth and obesity (Cupul-Uicab et al. 2013; Lamb et al. 2006; Mendez et al.

2011; Smink et al. 2008; Tang-Peronard et al. 2014; Valvi et al. 2014; Valvi et al. 2012; Verhulst et al. 2009; Warner et al. 2013; Warner et al. 2014). Prenatal DDT and DDE exposure has been positively associated with BMI in infancy (Mendez et al. 2011; Valvi et al. 2014; Verhulst et al. 2009) and later childhood (Tang-Peronard et al. 2014; Valvi et al. 2012; Warner et al. 2014). Findings for the association between prenatal PCB exposure and childhood obesity have been less consistent (Cupul-Uicab et al. 2013; Lamb et al. 2006; Mendez et al. 2011; Tang-Peronard et al. 2014; Valvi et al. 2014; Valvi et al. 2012; Verhulst et al. 2009). Prenatal HCB exposure has been positively associated with rapid growth in the first 6 months of life (Valvi et al. 2014) and obesity in infancy (Valvi et al. 2014) and childhood (Smink et al. 2008). Direct comparison of results from previous studies is limited by variations in exposure and outcome assessment and also because associations seem to be modified by other factors like sex, smoking or maternal BMI (Mendez et al. 2011; Tang-Peronard et al. 2014; Valvi et al. 2014; Valvi et al. 2012; Verhulst et al. 2009; Warner et al. 2014).

Cardiovascular risk factors measured in early childhood such as blood pressure and lipid levels are predictive of adulthood risk for coronary artery disease, suggesting that exposure to these factors early in life may induce changes in arteries that contribute to the development of atherosclerosis (Li et al. 2003; Raitakari et al. 2003). Exposure to environmental chemicals in adulthood has been associated with other cardiovascular traits such as higher blood pressure (La Merrill et al. 2013; Peters et al. 2014) and high serum lipid levels (Aminov et al. 2013; Patel et al. 2012) in adults. However, to our knowledge, there are no studies so far on the effect of prenatal POP exposure on offspring cardiovascular traits other than adiposity.

In the present study, we examined the association of maternal serum HCB, DDE and PCBs levels in pregnancy with offspring obesity and a range of cardiovascular risk factors (BMI, fat

mass, waist circumference, blood pressure, lipids, adiponectin, leptin, and C-reactive protein [CRP]) at 4 years of age in the Rhea mother-child cohort in Crete, Greece.

Materials and Methods

Study population

The Rhea study prospectively examines a population-based sample of pregnant women and their children at the prefecture of Heraklion, Crete, Greece. Methods are described in detail elsewhere (Chatzi et al. 2009). Briefly, female residents (Greek and immigrants) who became pregnant during a period of one year starting in February 2007 were contacted and asked to participate in the study. The first contact was made at the time of the first comprehensive ultrasound examination (mean \pm SD 11.96 \pm 1.49 weeks) and several contacts followed (6th month of pregnancy, at birth, 9 months, 1st year and 4 years after birth). To be eligible for inclusion in the study, women had to have a good understanding of the Greek language and be older than 16 years of age. Face-to-face structured questionnaires along with self-administered questionnaires and medical records were used to obtain information on several psychosocial, dietary, and environmental exposures during pregnancy and early childhood. The study was approved by the ethics committee of the University Hospital in Heraklion, Crete, Greece, and all participants provided written informed consent after complete description of the study.

1765 eligible women were approached during the enrollment period, 1610 (91%) agreed to participate and 1459 singletons (86%) were followed up until delivery. 1135 blood samples provided by the study participants were analyzed for POP exposure. 879 children participated at the 4 years follow up of the study, during which anthropometry and non-fasting blood samples were obtained from 785 children. From those, complete data for POP exposure, follow-up

interview and anthropometric measurements were available for 689 mother–child pairs and thus were eligible for analysis.

POP exposure

Maternal serum samples were collected at the first prenatal visit around the 3rd and 4th month of pregnancy, in 10 ml Silicone gel separator vacutainer (Becton Dickinson, UK), were centrifuged within 2 hours from blood collection at 2500rpm for 10min and were then stored in aliquots at -80°C until assayed. The POP analyses were performed in the National Institute for Health and Welfare, Chemical Exposure Unit, Kuopio, Finland with an Agilent 7000B gas chromatograph triple quadrupole mass spectrometer (GC-MS/MS). Pretreatment of serum samples for GC-MS/MS analysis has been described elsewhere (Koponen et al. 2013). Serum concentrations of six individual PCB congeners (IUPAC numbers: 118, 138, 153, 156, 170 and 180), HCB, DDT and DDE, and BDE-47 were determined. All the results were reported on whole weight and expressed in pg/ml serum, while samples bellow the limit of quantification (LOQ) were assigned the value $0.5 \times \text{LOQ}$. LOQ was 6 pg/ml for PCB118 and PCB 156; 10 pg/ml for HCB, DDE, PCB138, PCB153, PCB170, PCB180 and BDE47, and 50 pg/ml for DDT. We chose to use wet-weight levels for the POPs but adjusted for fasting maternal serum triglycerides [mean (\pm SD) = 128.4 (56.6) mg/dl] and cholesterol [208.9 (44.5) mg/dl] as continuous variables in all multivariable models to minimize potential biases associated with automatic lipid adjustment (Schisterman et al. 2005). The percentage of samples with levels of DDT above the LOQ was 35.3%. For PCB 156 and BDE-47, 53.9% and 23.1% were above the LOQ, respectively. Due to high percentages of samples below the LOQ, DDT and BDE-47 were not used in the statistical analyses. PCB 156 was included in analyses of summed PCB concentrations, but analyses were repeated after excluding this pollutant with no change in the results (data not shown).. POPs

were treated as continuous variables on a \log_{10} scale. We calculated total PCB concentrations by summing the concentrations of the 6 individual PCB congeners. We estimated associations using separate models for DDE, HCB, and the sum of PCBs, and using multipollutant models that included all three exposures.

Child anthropometry

Age- and sex-specific z-scores for weight at birth and at 6 months of age were calculated using internally generated growth reference curves. Rapid growth from birth to 6 months of age was defined as a z score weight gain greater than 0.67 standard deviation (SD) (Monteiro and Victora 2005). Children with a z score weight gain equal to or below 0.67 SD were characterized as slow/average growers and constituted the reference group.

At 4 years of age, body weight was measured once by a digital scale (SecaBellissima 841) to the nearest 0.1 kg with subjects standing without shoes and in light clothing. Height was measured to the nearest 0.1 cm with the use of a commercial stadiometer (Seca 213). The measurement of height was conducted without shoes and in a standing position with children keeping their shoulders in a relaxed position, their arms hanging freely and with their head aligned in Frankfurt horizontal plane. Child body mass index [BMI, $(\text{weight}/\text{height})^2$] was used to estimate age- and sex-specific z scores using the mean of the population under study. We categorized weight status at 4 years according to the International Obesity Task Force (IOTF) definitions (Cole and Lobstein 2012).

Waist circumference (WC) was measured in duplicate to the nearest 0.1 cm in the standing position, at the high point of the iliac crest at the end of a gentle expiration, using a flexible tape measure (Seca 201). Since there are no international WC percentile cut-off points defined to

identify central adiposity in children or adolescents, we decided to use a Greek national reference to define central adiposity as waist circumference (WC) \geq 90th percentile for age and sex (Linardakis et al. 2010).

Triceps, subscapular, suprailiac and quadriceps skinfold thicknesses were measured according to standardized techniques to the nearest 0.1mm using a calibrated Harpenden caliper (Harpenden HSK- BI, CE-0120). Three complete sets of measurement were taken consecutively and the mean was used as the representative value for each site. Fat mass was expressed as the sum of the four aforementioned skinfolds.

Child cardiometabolic risk factors

After 5 minutes rest in the seated position, trained research assistants measured systolic (SBP) and diastolic (DPB) blood pressure levels on the child's right arm right arm using an automatic oscillometric device (Dinamap, Pro Care 400) with a cuff of appropriate size for arm circumference. Five measurements were made with 1 minute intervals and the average of all measurements was used for analysis (Gillman and Cook 1995). Non-fasting blood samples were collected from the children at the end of the visit in 10 ml vacutainer tubes with the use of standard procedures, with samples immediately spun, separated, and frozen at -80°C . Analysis of lipids (total cholesterol and high-density lipoprotein cholesterol [HDL-C]) was performed by standard enzymatic methods (Medicon, Greece) on an automatic analyzer (AU5400 high-volume chemistry analyzer; Olympus America, Inc., Melville, New York). Leptin and adiponectin were measured by enzyme-linked immunosorbent assays (Leptin; DLP00, R&D Systems, Inc., MN and adiponectin; KHP0041, Invitrogen Corporation, CA) on an automatic analyzer (MAGO plus, Diamedix, Miami). C-reactive protein (CRP) levels were measured with a high-sensitivity homogenous immunoassay (ORS 6199, Beckman Coulter, USA) on an automatic analyzer

(AU5400 high-volume chemistry analyzer; Olympus America, Inc., Melville, New York) and abnormal levels were defined as >3 mg/L (Ford et al. 2005). All inter- and intra-assay coefficients of variation were < 5.5 %.

Statistical analysis

Descriptive analyses of the study population characteristics, exposures and outcomes were conducted. Generalized additive models (GAMs) were applied to explore the shape of the relationships between POPs in maternal serum and outcomes under study. These models indicated clear linear relationships (p -gain > 0.05). Multivariate generalized linear models for binary outcomes [with log link, Poisson distribution, and robust variance estimator (Zou 2004)] were used to estimate relative risks (RRs) and 95% confidence intervals (CIs) for the associations between the POP exposures and the outcome variables. Multivariate linear regression models were used to examine the association between POPs in maternal serum with continuous outcomes.

We used directed acyclic graphs (DAGs) to choose potential confounders for model adjustment. The set of variables selected for adjustment were maternal pre-pregnancy BMI (kg/m^2), maternal age at birth (years), parity (primiparous, multiparous), maternal educational level [low level: ≤ 9 years of mandatory schooling, medium level: > 9 years of schooling up to attending post-secondary school education and high level: attending university or having a university/technical college degree], smoking during pregnancy (never, ever), breastfeeding duration (months) and birth weight. This list of variables plus study child's sex, gestational age and exact age at outcome assessment, maternal serum levels of triglycerides and total cholesterol were the covariates selected *a priori* for model adjustment. Additionally, we assessed whether maternal weight before pregnancy (kg), gestational weight gain [GWG (kg)], ethnic origin (Greek, non-Greek),

residence (urban, rural), alcohol consumption during pregnancy (yes, no), delivery type (vaginal delivery, cesarean section), delivery hospital (private, public), marital status (married, not married) and working during pregnancy (yes, no) had further influence on the effect estimates. If inclusion of a variable altered the contaminant coefficient by 10% or more we retained the variable in the final set of covariates. The same set of covariates was used for all exposure–outcome combinations. None of the additionally tested variables were selected with this strategy. We looked for heterogeneity in associations related to child sex (male, female), maternal smoking (never, ever) and GWG categories [inadequate, adequate, excessive according to The Institute of Medicine (IOM) guidelines of 2009 based on pre-pregnancy BMI (IOM 2009)], or maternal pre-pregnancy BMI status ($<25 \text{ kg/m}^2$, $\geq 25 \text{ kg/m}^2$) by including interaction terms in the models for child z-BMI and by stratifying the sample. Statistical significance was defined by an alpha level of 0.10 for interaction terms and of 0.05 for all other effect estimates. We performed sensitivity analysis in order to explore remaining confounding. In particular, we repeated the analyses excluding children who had been born preterm (< 37 gestational weeks) or at low birth weight ($< 2500 \text{ g}$). Because child BMI could be an intermediate in the effect of prenatal POP exposure on waist circumference, leptin, blood pressure and lipids, we perform sensitivity analyses with adjustment for child BMI. Further, we repeated all analyses without adjustment for birth weight. Analyses were conducted using STATA software, version 13.0 (Statacorp, College Station, TX).

Results

Participating mothers were predominantly Greek, had a mean (\pm SD) age of 29.9 ± 5.0 years, about half of them had medium educational level (51%) and were multiparous (57%) (Table 1). Before pregnancy, 20% of mothers were overweight, almost all mothers (87%) initiated

breastfeeding and the mean length of breastfeeding was 4.1 ± 4.3 months. Table S1 in the Supplementary Material shows that mothers without offspring follow-up data were more likely to be younger, less educated and not Greek.

About half of the children (52%) were boys and 49% were vaginally delivered (Table 1). The prevalence of rapid growth in the first 6 months of life was 35% (N=203). Rapid growers were at higher risk for overweight/obesity at 4 years of age (adj. RR= 1.94; 95% CI: 1.46, 2.56). At the 4-year follow-up, the mean (\pm SD) offspring BMI was 16.4 ± 1.9 kg/m². In total, 99 (14%) children were classified as overweight and an additional 48 (7%) were obese at 4 years of age. Mean waist circumference was 53.6 ± 4.7 cm and a total of 80 (12%) children had a waist circumference \geq 90th percentile for age.

Maternal POP concentrations are presented in Table 2. The highest concentrations were found for DDE, followed by total PCBs, HCB, and DDT. Spearman correlation coefficients (p-value) were 0.48 (<0.001) for DDE-PCBs, 0.50 (<0.001) for DDE-HCB, and 0.64 (<0.001) for PCBs-HCB. Mothers that were older, multiparous, and had higher pre-pregnancy BMI, had higher mean POP levels in early pregnancy (see Supplemental Material, Table S2). Prenatal DDE and HCB levels were similar between women whose offspring had outcome measurements and those that did not, while PCB levels were lower in women without offspring follow-up data (see Supplemental Material, Table S1). GAMs examining the shape of the relationships of HCB, DDE and PCBs with child z-BMI at 4 years showed no significant departures from linearity overall and separately in boys and girls (Figure 1).

Table 3 shows adjusted results for maternal POP levels in relation to offspring obesity-related outcomes. Because models were run with POPs on the log₁₀ scale, each unit of increase means a

10-fold increase in the concentration of each contaminant. Prenatal HCB concentrations were positively but not significantly associated with rapid growth in the first 6 months (adj. RR=1.94; 95% CI: 0.99, 3.77). At 4 years of age, prenatal HCB concentration was associated with about ½ standard deviation higher BMI z score (adj. β = 0.49; 95% CI: 0.12, 0.86), and a higher prevalence of obesity (adj. RR=8.14; 95% CI: 1.85, 35.81); however, given the small number of children classified as obese (n=32, 6%), the estimates were imprecise as shown by the wide confidence intervals. Increased *in utero* exposure to HCB was also significantly associated with WC \geq 90th percentile (adj. RR=3.49; 95% CI: 1.08, 11.28), and with a higher sum of skinfold thickness (adj. β = 7.71 mm; 95% CI: 2.04, 13.39) at 4 years of age. DDE was also associated with an increase in the BMI z score (adj. β = 0.27; 95% CI: 0.04, 0.5) and higher prevalence of abdominal obesity (adj. RR=3.76; 95% CI: 1.70, 8.30) at 4 years of age. In general, PCBs were positively but not significantly associated with offspring obesity outcomes (Table 3).

Table 4 presents the associations between maternal POP exposure in pregnancy and offspring cardiometabolic risk factors at 4 years of age. Higher systolic blood pressure was observed for increasing levels of HCB (adj. β = 4.34 mmHg; 95% CI: 0.63, 8.05), while increasing levels of DDE were associated with higher diastolic blood pressure (adj. β = 1.79 mmHg; 95% CI: 0.13, 3.46). HCB was also associated with DBP and DDE was also associated with SBP, though the observed associations were not significant. Maternal HCB and DDE serum levels were positively associated with child blood leptin concentrations at 4 years of age (adj. β = 2.15 ng/ml; 95% CI: 0.42, 3.89; adj. β = 1.21 ng/ml; 95% CI: 0.16, 2.27 respectively). Associations attenuated and were not anymore significant when we adjusted for child BMI at age of outcome assessment (adj. β = 1.01 ng/ml; 95% CI: -0.21, 2.23; adj. β = 0.47 ng/ml; 95% CI: -0.28, 1.21 respectively). Maternal HCB, DDE and PCBs levels were positively associated with elevated offspring CRP

levels at 4 years of age, although the observed associations did not reach statistical significance (Table 4). However, only a small number of children had abnormal CRP levels (n=46, 11%). Child lipids (total cholesterol and HDL) and adiponectin levels were not associated with any of the measured POPs (see Supplemental Material, Table S3). Estimates from multi-pollutant model including DDE, HCB and PCBs indicate stronger associations with HCB when adjusted for the other two exposures, and comparisons with single-pollutant model estimates suggest that associations with PCBs were subject to the greatest amount of positive confounding by the other exposures (see Supplemental Material, Table S4).

We found no evidence of effect modification for BMI z score at 4 years (see Supplemental Material, Table S4), or any of the other outcomes (data not shown) by child sex, smoking status during pregnancy, pre-pregnancy BMI or weight gain during pregnancy (all p-interaction > 0.10). In order to evaluate the influence of breastfeeding on rapid growth, we performed a stratified analysis by breastfeeding duration which showed no substantial differences in observed estimates (data not shown). Sensitivity analyses excluding preterm newborns (<37 gestational weeks; n=66-45) and infants with low birth weight (i.e., <2500 g, n=23-18) did not meaningfully change our results (see Supplemental Material, Table S5). Estimates were also not meaningfully changed by the exclusion of birth weight from the models. Inclusion of child BMI at 4 years in the models for waist circumference slightly decreased the effect estimates, but did not influence their statistical significance (data not shown). Finally, adjusting models of associations for blood pressure and lipids by child BMI at 4 years did not change the results substantially (data not shown).

Discussion

In this population-based birth cohort study, *in utero* exposure to HCB and DDE was associated with childhood obesity and higher offspring blood pressure levels at 4 years. To our knowledge this is the first study examining the relationship between prenatal exposure to organochlorine chemicals and child cardiometabolic risk factors other than obesity.

We found that higher prenatal HCB and DDE concentrations were significantly associated with increased BMI z scores, and adiposity at 4 years of age. Our findings for HCB are consistent with previous studies that reported positive associations with offspring obesity at 14 months (Valvi et al. 2014) and 6 years (Smink et al. 2008). On the contrary, Delvaux et al. (2014) reported no significant association of HCB measured in cord blood and markers of abdominal obesity at 7 to 9 years. Low-level prenatal DDE exposure has also been linked to elevated BMI at 14 months of age (Mendez et al. 2011; Valvi et al. 2014), at 3 years of age (Verhulst et al. 2009), at 6.5 (Valvi et al. 2012), 9 years of age (Warner et al. 2014), and markers of abdominal fat in girls at 7 to 9 years (Delvaux et al. 2014). However, recent studies in populations with higher exposures reported null associations between prenatal DDE exposure and BMI at 12 months (Garced et al. 2012), 18 months (Cupul-Uicab et al. 2010) and 7 years of age (Cupul-Uicab et al. 2013; Warner et al. 2013) suggesting a non-monotonic dose response relationship (Vandenberg et al. 2012). Potential effect modification by sex and other factors found in previous studies (Cupul-Uicab et al. 2013; Mendez et al. 2011; Tang-Peronard et al. 2014; Valvi et al. 2012; Warner et al. 2014) was not supported in our study maybe due to smaller population subgroups and should be further explored in larger populations. We did not find an association of prenatal PCBs exposure with child adiposity, in line with other studies reporting null associations between prenatal PCB exposure and elevated BMI (Cupul-Uicab et al. 2013;

Delvaux et al. 2014; Mendez et al. 2011; Valvi et al. 2014). Although Verhulst et al. (2009) reported a very small increase in BMI z-scores associated with PCB exposure in children 1–3 years old and Tang-Péronard et. al (2014) an association with BMI for 7-year old girls, the preponderance of evidence does not suggest a strong relationship between prenatal PCB exposure and later adiposity. Nevertheless, in this study, estimates from multi-pollutant models suggest that associations with PCBs might be positively confounded by other POPs exposures.

The mechanisms by which prenatal exposure to POPs, such DDE and HCB might be related to increased body fatness are still not clear (Tang-Peronard et al. 2011). In vitro studies confirm that DDE exposure at environmentally relevant levels increases proliferation (Chapados et al. 2012) and differentiation (Moreno-Aliaga and Matsumura 2002) of preadipocytes. First, it is possible that obesogenic effects of POPs are mediated by sex steroid dysregulation. DDE is both an estrogen receptor agonist and androgen receptor antagonist, while HCB has been suggested to be an androgen receptor and an estrogen-related receptor antagonist (Li et al. 2008). Knockout models of sex steroid pathway components, confirm that sex steroids are involved in adipocyte hypertrophy and hyperplasia (Heine et al. 2000; Murata et al. 2002). Secondly, DDT has been associated with increased mRNA expression of the peroxisome proliferator activated receptor γ (PPAR γ) (Moreno-Aliaga and Matsumura 2002) which constitutes a major regulator of adipogenesis; it is primarily expressed in adipose tissue, and its activation promotes adipocyte differentiation as well as the induction of lipogenic enzymes (Diamanti-Kandarakis et al. 2009). Third, it is possible that organochlorine compounds mediate their obesogenic effect through central integration of energy balance (Grun and Blumberg 2009). We found a positive association between prenatal HCB and DDE levels and child leptin levels, a hormone heavily involved in the control of energy balance (Yu et al. 1997). Consistent with our results, in vitro

studies have shown that exposure of mature adipocytes to DDE resulted in increased leptin release (Howell and Mangum 2011). Further studies are needed to better understand how prenatal and early life exposure to obesogenic chemicals, such as organochlorines, can program exposed individuals to gain weight and what role modulation of hormone receptor expression or activity early in life plays in this process.

Prenatal HCB and DDE levels were associated with higher child blood pressure at preschool age. Exposure to environmental chemicals, such as PCBs and DDE, has been associated with higher blood pressure (Goncharov et al. 2011; La Merrill et al. 2013; Lind et al. 2014; Peters et al. 2014) in adults. An association between exposure to HCB and increased heart weight has been reported in mice (Arnold et al. 1986), while exposure to other pollutants like dioxins and PCBs has also been associated with hypertension in animal studies (Kopf et al. 2008; Lind et al. 2004). Despite the evidence showing an association with cardiovascular parameters, animal models elucidating the mechanisms underlying these effects are limited.

The POPs levels in the study population were close to but generally lower than those of other pregnant populations. Exposure to HCB (median 0.08 ug/L) was much lower compared to the median exposure in other studies [Valvi et al. (2012) 0.67 ug/L, Cupul-Uicab et al. (2013) 0.24 ug/L, and Smink et al. (2008) 0.68 ug/L]. Similarly DDE (1.98 ug/L) levels were much lower compared to those reported by Cupul-Uicab et al. (2013) in the United States (24.59 ug/L) but somewhat higher than another European study (1.06 ug/L) published by Valvi et al. (2012). PCBs were also lower (0.32 ug/l) than previously reported levels in pregnant women [Valvi et al. (2012) 0.75 ug/L, Cupul-Uicab et al. (2013) 2.74 ug/L]. Therefore, it is possible that relations observed within our study population estimate the association of these chemicals with child health outcomes only within low exposure levels.

Strengths of the present study include the population-based prospective design, the relatively large sample size compared to previous studies and the detailed childhood body fat and cardiometabolic measurements. The fairly homogenous population with regard to factors such as diet, breastfeeding, country of origin, and socioeconomic status can reduce uncontrolled confounding. The levels of attrition in the Rhea cohort are similar to those found in other birth cohort studies (Guxens et al. 2012; Jaddoe et al. 2012), while we found no evidence of differences in prenatal HCB and DDE levels between participants and non-participants. We were able to measure levels of several POPs in first trimester maternal serum using state-of-the-art laboratory techniques and we were able to assess the adverse effect of combined exposure to several contaminants. However, our study has also some limitations. We could not separate the potential effects of postnatal exposures since we have not measured POPs at other time points, but were able to control for these possible effects by taking into account breastfeeding duration. We cannot also rule out the possibility that prenatal and/or postnatal exposure to other unmeasured chemicals correlated to POPs may have confounded the associations under study. Although we studied a broad spectrum of outcomes we did not do multiple comparisons as some of the outcomes examined are inter-correlated (adiposity measurements); therefore we did not apply Bonferroni type corrections. Finally, although we incorporated extensive information on potential social and environmental factors that are associated with childhood adiposity and cardiometabolic risk factors, we acknowledge that residual confounding from unmeasured covariates such as parental income or social class is still possible.

Conclusions

This study suggests that prenatal exposure to low levels of DDE and HCB may be associated with greater offspring adiposity and higher blood pressure levels in early childhood. These

findings have important public health implications because of the high and increasing prevalence of childhood obesity and ubiquity of POPs exposure. Our future research in the Rhea cohort will examine whether these compounds are associated with alterations in growth at late childhood and puberty onset.

References

- Aminov Z, Haase RF, Pavuk M, Carpenter DO. 2013. Analysis of the effects of exposure to polychlorinated biphenyls and chlorinated pesticides on serum lipid levels in residents of Anniston, Alabama. *Environ Health* 12:108.
- Arnold DL, Moodie CA, Collins BT, Zawidzka ZZ, Krewski DR. 1986. Two-generation chronic toxicity study with hexachlorobenzene in the rat. *IARC Sci Publ*(77):405–410.
- Baillie-Hamilton PF. 2002. Chemical toxins: a hypothesis to explain the global obesity epidemic. *J Altern Complement Med* 8(2):185–192.
- Chapados NA, Casimiro C, Robidoux MA, Haman F, Batal M, Blais JM, et al. 2012. Increased proliferative effect of organochlorine compounds on human preadipocytes. *Mol Cell Biochem* 365(1-2):275–278.
- Chatzi L, Plana E, Daraki V, Karakosta P, Alekakis D, Tsatsanis C, et al. 2009. Metabolic syndrome in early pregnancy and risk of preterm birth. *Am J Epidemiol* 170(7):829–836.
- Cole TJ, Lobstein T. 2012. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 7(4):284–294.
- Cupul-Uicab LA, Hernandez-Avila M, Terrazas-Medina EA, Pennell ML, Longnecker MP. 2010. Prenatal exposure to the major DDT metabolite 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene (DDE) and growth in boys from Mexico. *Environ Res* 110(6):595–603.
- Cupul-Uicab LA, Klebanoff MA, Brock JW, Longnecker MP. 2013. Prenatal exposure to persistent organochlorines and childhood obesity in the US collaborative perinatal project. *Environ Health Perspect* 121(9):1103–1109.
- Delvaux I, Van Cauwenberghe J, Den Hond E, Schoeters G, Govarts E, Nelen V, et al. 2014. Prenatal exposure to environmental contaminants and body composition at age 7-9 years. *Environ Res* 132:24–32.
- Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, et al. 2009. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30(4):293–342.
- Ford ES, Ajani UA, Mokdad AH. 2005. The metabolic syndrome and concentrations of C-reactive protein among U.S. youth. *Diabetes Care* 28(4):878–881.

- Garced S, Torres-Sanchez L, Cebrian ME, Claudio L, Lopez-Carrillo L. 2012. Prenatal dichlorodiphenyldichloroethylene (DDE) exposure and child growth during the first year of life. *Environ Res* 113:58–62.
- Gillman MW, Cook NR. 1995. Blood pressure measurement in childhood epidemiological studies. *Circulation* 92(4):1049–1057.
- Goncharov A, Pavuk M, Foushee HR, Carpenter DO. 2011. Blood pressure in relation to concentrations of PCB congeners and chlorinated pesticides. *Environ Health Perspect* 119(3):319–325.
- Grun F, Blumberg B. 2006. Environmental obesogens: organotins and endocrine disruption via nuclear receptor signaling. *Endocrinology* 147(6 Suppl):S50–55.
- Grun F, Blumberg B. 2009. Endocrine disrupters as obesogens. *Mol Cell Endocrinol* 304(1-2):19–29.
- Guxens M, Ballester F, Espada M, Fernandez MF, Grimalt JO, Ibarluzea J, et al. 2012. Cohort Profile: the INMA--Infancia y Medio Ambiente--(Environment and Childhood) Project. *Int J Epidemiol* 41(4):930–940.
- Heine PA, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS. 2000. Increased adipose tissue in male and female estrogen receptor-alpha knockout mice. *Proc Natl Acad Sci U S A* 97(23):12729–12734.
- Howell G, 3rd, Mangum L. 2011. Exposure to bioaccumulative organochlorine compounds alters adipogenesis, fatty acid uptake, and adipokine production in NIH3T3-L1 cells. *Toxicol In Vitro* 25(1):394–402.
- Jaddoe VW, van Duijn CM, Franco OH, van der Heijden AJ, van Iizendoorn MH, de Jongste JC, et al. 2012. The Generation R Study: design and cohort update 2012. *Eur J Epidemiol* 27(9):739–756.
- Kopf PG, Huwe JK, Walker MK. 2008. Hypertension, cardiac hypertrophy, and impaired vascular relaxation induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin are associated with increased superoxide. *Cardiovasc Toxicol* 8(4):181–193.
- Koponen J, Rantakokko P, Airaksinen R, Kiviranta H. 2013. Determination of selected perfluorinated alkyl acids and persistent organic pollutants from a small volume human serum sample relevant for epidemiological studies. *J Chromatogr A* 1309:48–55.

- La Merrill M, Birnbaum LS. 2011. Childhood obesity and environmental chemicals. *Mt Sinai J Med* 78(1):22–48.
- La Merrill M, Cirillo PM, Terry MB, Krigbaum NY, Flom JD, Cohn BA. 2013. Prenatal exposure to the pesticide DDT and hypertension diagnosed in women before age 50: a longitudinal birth cohort study. *Environ Health Perspect* 121(5):594–599.
- Lamb MR, Taylor S, Liu X, Wolff MS, Borrell L, Matte TD, et al. 2006. Prenatal exposure to polychlorinated biphenyls and postnatal growth: a structural analysis. *Environ Health Perspect* 114(5):779–785.
- Li J, Li N, Ma M, Giesy JP, Wang Z. 2008. In vitro profiling of the endocrine disrupting potency of organochlorine pesticides. *Toxicol Lett* 183(1-3):65–71.
- Li S, Chen W, Srinivasan SR, Bond MG, Tang R, Urbina EM, et al. 2003. Childhood cardiovascular risk factors and carotid vascular changes in adulthood: the Bogalusa Heart Study. *JAMA* 290(17):2271–2276.
- Linardakis M, Sarri K, Bertsiias G, Papadaki A, Kafatos A. 2010. Waist circumference and waist-to-height percentiles for the youth of Crete, Greece. *Int J Child Health and Hum Dev* 3:81–93.
- Lind PM, Orberg J, Edlund UB, Sjoblom L, Lind L. 2004. The dioxin-like pollutant PCB 126 (3,3',4,4',5-pentachlorobiphenyl) affects risk factors for cardiovascular disease in female rats. *Toxicol Lett* 150(3): 293–299.
- Lind PM, Penell J, Salihovic S, van Bavel B, Lind L. 2014. Circulating levels of p,p'-DDE are related to prevalent hypertension in the elderly. *Environ Res* 129:27–31.
- Mendez MA, Garcia-Esteban R, Guxens M, Vrijheid M, Kogevinas M, Goni F, et al. 2011. Prenatal organochlorine compound exposure, rapid weight gain, and overweight in infancy. *Environ Health Perspect* 119(2):272–278.
- Monteiro PO, Victora CG. 2005. Rapid growth in infancy and childhood and obesity in later life—a systematic review. *Obes Rev* 6(2):143–154.
- Moreno-Aliaga MJ, Matsumura F. 2002. Effects of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane (p,p'-DDT) on 3T3-L1 and 3T3-F442A adipocyte differentiation. *Biochem Pharmacol* 63(5):997–1007.
- Murata Y, Robertson KM, Jones ME, Simpson ER. 2002. Effect of estrogen deficiency in the male: the ArKO mouse model. *Mol Cell Endocrinol* 193(1-2):7–12.

- Needham LL, Grandjean P, Heinzow B, Jorgensen PJ, Nielsen F, Patterson DG, Jr., et al. 2011. Partition of environmental chemicals between maternal and fetal blood and tissues. *Environ Sci Technol* 45(3):1121–1126.
- Patel CJ, Cullen MR, Ioannidis JP, Butte AJ. 2012. Systematic evaluation of environmental factors: persistent pollutants and nutrients correlated with serum lipid levels. *Int J Epidemiol* 41(3):828–843.
- Peters JL, Patricia Fabian M, Levy JI. 2014. Combined impact of lead, cadmium, polychlorinated biphenyls and non-chemical risk factors on blood pressure in NHANES. *Environ Res* 132C:93–99.
- Porta M, Puigdomenech E, Ballester F, Selva J, Ribas-Fito N, Llop S, et al. 2008. Monitoring concentrations of persistent organic pollutants in the general population: the international experience. *Environ Int* 34(4):546–561.
- Raitakari OT, Juonala M, Kahonen M, Taittonen L, Laitinen T, Maki-Torkko N, et al. 2003. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *JAMA* 290(17):2277–2283.
- Rasmussen K.M., Yaktine A.L. (Eds.), *Weight gain during pregnancy: reexamining the guidelines*, National Academies Press, Washington, DC (2009)
- Schisterman EF, Whitcomb BW, Louis GM, Louis TA. 2005. Lipid adjustment in the analysis of environmental contaminants and human health risks. *Environ Health Perspect* 113(7):853–857.
- Smink A, Ribas-Fito N, Garcia R, Torrent M, Mendez MA, Grimalt JO, et al. 2008. Exposure to hexachlorobenzene during pregnancy increases the risk of overweight in children aged 6 years. *Acta Paediatr* 97(10):1465–1469.
- Stockholm Convention on Persistent Organic Pollutants. 2009. Available: <http://chm.pops.int> [accessed 10 November 2014].
- Tang-Peronard JL, Andersen HR, Jensen TK, Heitmann BL. 2011. Endocrine-disrupting chemicals and obesity development in humans: a review. *Obes Rev* 12(8):622–636.
- Tang-Peronard JL, Heitmann BL, Andersen HR, Steuerwald U, Grandjean P, Weihe P, et al. 2014. Association between prenatal polychlorinated biphenyl exposure and obesity development at ages 5 and 7 y: a prospective cohort study of 656 children from the Faroe Islands. *Am J Clin Nutr* 99(1):5–13.

- Valvi D, Mendez MA, Garcia-Esteban R, Ballester F, Ibarluzea J, Goni F, et al. 2014. Prenatal exposure to persistent organic pollutants and rapid weight gain and overweight in infancy. *Obesity (Silver Spring)* 22(2):488–496.
- Valvi D, Mendez MA, Martinez D, Grimalt JO, Torrent M, Sunyer J, et al. 2012. Prenatal concentrations of polychlorinated biphenyls, DDE, and DDT and overweight in children: a prospective birth cohort study. *Environ Health Perspect* 120(3):451–457.
- Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs DR, Jr., Lee DH, et al. 2012. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocr Rev* 33(3):378–455.
- Verhulst SL, Nelen V, Hond ED, Koppen G, Beunckens C, Vael C, et al. 2009. Intrauterine exposure to environmental pollutants and body mass index during the first 3 years of life. *Environ Health Perspect* 117(1):122–126.
- Warner M, Aguilar Schall R, Harley KG, Bradman A, Barr D, Eskenazi B. 2013. In utero DDT and DDE exposure and obesity status of 7-year-old Mexican-American children in the CHAMACOS cohort. *Environ Health Perspect* 121(5):631–636.
- Warner M, Wesselink A, Harley KG, Bradman A, Kogut K, Eskenazi B. 2014. Prenatal Exposure to Dichlorodiphenyltrichloroethane and Obesity at 9 Years of Age in the CHAMACOS Study Cohort. *Am J Epidemiol* 179(11):1312–1322.
- WHO. 2010. Persistent Organic Pollutants: Impact on Child Health. Available: http://www.who.int/ceh/publications/persistent_organic_pollutant/en/index.html [accessed 4 June 2014].
- Yu WH, Kimura M, Walczewska A, Karanth S, McCann SM. 1997. Role of leptin in hypothalamic-pituitary function. *Proc Natl Acad Sci U S A* 94(3):1023–1028.
- Zou G. 2004. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 159(7):702–706.

Table 1. Maternal and child characteristics, Mother-child cohort “Rhea” in Crete, Greece (n=689).

Characteristic	N	% or Mean ± SD
Maternal characteristics		
Maternal age (years)	681	29.9±5.0
<20	18	2.6
≥20-30	276	40.5
≥30-40	368	54.1
≥40	19	2.8
Pre-pregnancy BMI (kg/m ²)	675	24.4±4.7
Underweight (<18.5)	22	3.3
Normal (≥18.5-25)	437	64.7
Overweight (≥25-30)	137	20.3
Obese (≥30)	79	11.7
Weight gain during pregnancy (kg)	566	13.8±5.6
Ethnic origin		
Other	37	5.4
Greek	648	94.6
Education		
Low	105	15.5
Medium	347	51.2
High	226	33.3
Parity		
Primiparous	284	42.9
Multiparous	378	57.1
Smoking during pregnancy		
Never	432	64.7
Ever	236	35.3
Type of delivery		
Vaginal	339	49.4
Caesarian	347	50.6
Breastfeeding (months)		
Never	85	12.8
Ever	578	87.2
BMI at child’s 4 years		
Underweight (<18.5)	20	3.0
Normal (≥18.5-25)	337	50.9
Overweight (≥25-30)	201	30.4
Obese (≥30)	104	15.7
Child characteristics at birth		
Sex		
Boy	358	52.0
Girl	331	48.0

Characteristic	N	% or Mean ± SD
Birth weight (g)	686	3205±451.7
Gestational age (completed weeks)	686	38.2±1.5
Preterm (<37 weeks)		
Yes	84	12.2
No	602	87.8
Low Birth Weight		
Yes	30	4.4
No	656	95.6
Rapid growth		
Yes	203	35.2
No	374	64.8
Child characteristics at 4 years		
Weight at 4 years (kg)	689	18.2±3.0
Height (cm)	689	105.2
BMI (kg/m ²)	689	16.4±1.9
Underweight	15	2.2
Normal	527	76.5
Overweight	99	14.4
Obese	48	6.9
Triceps skinfold thickness (mm)	657	10.33±2.77
Subscapular skinfold thickness (mm)	661	6.93±2.61
Suprailiac skinfold thickness (mm)	643	7.90±3.86
Quadiceps skinfold thickness (mm)	621	15.54±5.40
Sum of skinfolds (mm)	668	39.1±14.2
Waist circumference (cm)	683	53.6±4.7
Systolic blood pressure (mmHg)	552	90.3±8.1
Diastolic blood pressure (mmHg)	552	53.6±5.4
Total Cholesterols (mg/dl)	588	157.1±28.6
HDL Cholesterol (mg/dl)	588	47.1±10.6
LDL Cholesterols (mg/dl)	588	96.6±25.5
C-reactive protein (ml/dl)	540	0.2±0.4
Leptin (ng/ml)	580	2.9±3.6
Adiponectin (ul/ml)	581	15.4±8.7

Table 2. First trimester maternal serum POP levels (pg/ml, n=689), Mother-child cohort “Rhea” in Crete, Greece (n=689).

Contaminants	GM (95% CI)			Percentile			LOQ	% above LOQ
		min	max	25th	50th	75th		
HCB	89.4 (85.6, 93.2)	19.5	1330.5	61.7	82.5	116.7	10	100
DDE	2036.2 (1910.2, 2170.5)	181.6	23175.4	1187.6	1981.2	3514.1	10	100
PCB118	17.1 (16.4, 17.9)	3 ^a	143.6	12.0	17.7	25.1	6	96.8
PCB153	124.4 (119.0, 130.2)	13.0	1348.9	85.5	125.2	188.1	10	100
PCB138	66.5 (63.5, 69.6)	5 ^a	742.8	45.3	67.8	102.0	10	99.7
PCB156	6.0 (5.7, 6.4)	3 ^a	81.5	3 ^a	6.4	10.7	6	53.9
PCB180	67.0 (63.7, 70.5)	5 ^a	979.6	44.1	66.3	104.5	10	99.7
PCB170	33.4 (31.6, 35.2)	5 ^a	542.3	22.0	33.7	53.4	10	96.2
Total PCBs	319.1 (304.9, 333.9)	34.0	3758.3	216.7	319.4	483.1		

^aValue is LOQ/2.

Table 3. Associations of *in utero* POP exposures with early rapid growth and obesity related outcomes at 4 years of age, Mother-child cohort Rhea in Crete, Greece.

Outcome	Cases (%) / total	Exposure ^a	Adjusted Model ^b
Rapid Growth 0-6 months ^c [RR (95% CI)]	157 (34.3)/458	HCB	1.94 (0.99, 3.77)
		DDE	1.33 (0.89, 1.99)
		PCBs	1.20 (0.59, 2.46)
BMI z-score at 4 years [β (95% CI)]	531	HCB	0.49 (0.12, 0.86)
		DDE	0.27 (0.04, 0.51)
		PCBs	0.21 (-0.20, 0.63)
Obesity at 4 years [RR (95% CI)]	32 (6.0)/531	HCB	8.14 (1.85, 35.81)
		DDE	3.80 (1.19, 12.14)
		PCBs	3.91 (0.49, 31.27)
WC \geq 90th percentile ^d [RR (95% CI)]	56 (10.7)/526	HCB	3.49 (1.08, 11.28)
		DDE	3.76 (1.70, 8.30)
		PCBs	2.08 (0.65, 6.71)
Sum of skinfolds (mm) at 4 years [β (95% CI)]	515	HCB	7.71 (2.04, 13.39)
		DDE	2.75 (-0.86, 6.35)
		PCBs	5.74 (-0.68, 12.16)

^aPOP concentrations are log₁₀ transformed, in pg/ml. ^bAdjusted for maternal serum triglycerides and cholesterol, maternal age, pre-pregnancy BMI, parity, maternal educational level, smoking status during pregnancy, breastfeeding duration, child sex, birth weight, gestational age and exact age at 4 year examination. ^cRapid growth models are adjusted for maternal serum triglycerides and cholesterol, maternal age, pre-pregnancy BMI, parity, maternal educational level, smoking status during pregnancy, breastfeeding duration, child sex and gestational age.

^dWC \geq 58.6cm

Table 4. Associations of *in utero* POPs exposure with cardiometabolic risk factors at 4 years of age, Mother-child cohort “Rhea” in Crete, Greece.


Outcome	Cases (%) / total	Exposure ^a	Adjusted Model ^b
Systolic blood pressure (mmHg) [β (95% CI)]	427	HCb	4.34 (0.63, 8.05)
		DDE	2.31 (-0.07, 4.69)
		PCBs	2.16 (-2.03, 6.34)
Diastolic blood pressure (mmHg) [β (95% CI)]	427	HCb	2.48 (-0.13, 5.09)
		DDE	1.79 (0.13, 3.46)
		PCBs	-0.49 (-3.43, 2.45)
C-reactive protein >3 mg/L [RR (95% CI)]	46 (11.3)/409	HCb	2.88 (0.86, 9.64)
		DDE	2.23 (0.94, 5.29)
		PCBs	4.50 (0.89, 22.76)
Leptin (ng/ml) [β (95% CI)]	440	HCb	2.15 (0.42, 3.89)
		DDE	1.21 (0.16, 2.27)
		PCBs	1.55 (-0.42, 3.52)

^aPOP concentrations are \log_{10} transformed, in pg/ml. ^bAdjusted for maternal serum triglycerides and cholesterol, maternal age, pre-pregnancy BMI, parity, maternal educational level, smoking status during pregnancy, breastfeeding duration, child sex, birth weight, gestational age and exact age at 4 year examination.

Figure Legend

Figure 1. GAMS; adjusted associations (95% CIs) of HCB overall (A) and in boys (B) and girls (C), DDE overall (D) and in boys (E) and girls (F) and PCBs overall (D) and in boys (E) and girls (F) with BMI z score at 4 years. Adjusted for maternal serum triglycerides and cholesterol, maternal age, pre-pregnancy BMI, parity, maternal educational level, smoking status during pregnancy, breastfeeding duration, child sex, birth weight, gestational age and exact age at 4 year examination. ++, observations.

Figure 1.



4