

Association of residential air pollution, noise, and greenspace with initial ischemic stroke severity.

Rosa Maria Vivanco-Hidalgo MD PhD MPH.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain, rvivanco@imim.es

Carla Avellaneda-Gómez MD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. ^bUniversitat Autònoma de Barcelona (UAB), Barcelona, Spain. cavellaneda@parcdesalutmar.cat

Payam Dadvand MD PhD.

^cBarcelonaInstitute for Global Health (ISGlobal), C/DrAiguader 88, CP08003, Barcelona, Spain; ^dUniversitat PompeuFabra (UPF), Barcelona, Spain; CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain, payam.dadvand@isglobal.org

Marta Cirach

^cISGlobal, C/DrAiguader 88, CP08003, Barcelona. marta.cirach@isglobal.org

ÁngelOis MD PhD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. aois@parcdesalutmar.cat

Alejandra Gómez González MD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. 60558@parcdesalutmar.cat

Ana Rodriguez-Campello MD PhD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. 92269@parcdesalutmar.cat

Pablo de Ceballos, MD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain PDeCeballos@parcdesalutmar.cat

Xavier Basagaña, PhD.

^cBarcelona Institute for Global Health (ISGlobal), Barcelona, Spain;

^dUniversitat Pompeu Fabra (UPF), Barcelona, Spain; CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain. xavier.basagana@isglobal.org

Ana Zabalza, MD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain anazabalza@gmail.com

Elisa Cuadrado-Godia MD PhD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. ecuadrado@parcdesalutmar.cat

Jordi Sunyer MD PhD.

^cBarcelona Institute for Global Health (ISGlobal), C/DrAiguader 88, CP08003, Barcelona, Spain;^dUniversitat Pompeu Fabra (UPF), Barcelona, Spain; CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain. jordi.sunyer@isglobal.org

Jaume Roquer MD PhD.

^aHospital del Mar Medical Research Institute, C/DrAiguader 88, CP08003, Barcelona, Spain. jroquer@parcdesalutmar.cat

Gregory A. Wellenius ScD.

^eDepartment of Epidemiology, Brown University School of Public Health, Providence, RI 02912, USA. gregory_wellenius@brown.edu

Corresponding authors

Rosa Maria Vivanco-Hidalgo, rvivanco@imim.es. C/ Doctor Aiguader 88, Barcelona, Spain. Tel: 0034933160611

Jaume Roquer, jroquer@parcdesalutmar.cat C/ Doctor Aiguader 88, Barcelona, Spain. Tel: 0034 933160611

Cover title: Environmental determinants of ischemic stroke severity

Tables 5, Figures 1

Word count: 4286

Abstract:

Background and Purpose: A number of environmental risk factors of acute ischemic stroke have been identified, but few studies have evaluated the influence of the outdoor environment on stroke severity. We assessed the association of residential ambient fine particulate matter air pollution (PM_{2.5}), noise, and surrounding greenspace with initial stroke severity.

Methods: We obtained data on patients hospitalized with acute ischemic stroke from a hospital-based prospective stroke register (2005-2014) in Barcelona. We estimated residential PM_{2.5} based on an established land use regression model, greenspace as the average satellite-based Normalized Difference Vegetation Index (NDVI) within a 300 m buffer of the residence, and daily (*L_{day}*), evening (*L_{evening}*), night (*L_{night}*) and average noise (*L_{den}*) level at the street nearest to the residential address using municipal noise models. Stroke severity was assessed at the time of hospital presentation using the National Institute of Health Stroke Scale (NIHSS). We used logistic regression and binomial models to evaluate the associations of PM_{2.5}, greenspace, and noise with initial stroke severity adjusting for potential confounders.

Results : Among 2761 patients, higher residential surrounding greenspace was associated with lower risk of severe stroke (OR for NIHSS>5, 0.75; 95% CI: 0.60-0.95), while, living in areas with higher *L_{den}* was associated with a higher risk of severe stroke (OR, 1.30; 95% CI: 1.02-1.65). PM_{2.5} was not associated with initial stroke severity.

Conclusions: In an urban setting, surrounding greenspace and traffic noise at home are associated with initial stroke severity, suggesting an important influence of the built environment on the global burden of ischemic stroke.

Key words: ischemic stroke air pollution noise greenspace

Source of funding

The study was funded by the Spanish Ministry of Economy through the Carlos III Health Institute (ISCIII-FISFEDER-ERDFPI15/00445), the Spanish Ministry of Health (Ministerio de Sanidad y Consumo, Instituto de Salud Carlos III FEDER, RD16/0019/0002. INVICTUS-PLUS), and grant number R01-ES020871 from the National Institute of Environmental Health Science (NIEHS, NIH). Payam Dadvand is funded by a Ramón y Cajal fellowship (RYC-2012-10995) awarded by the Spanish Ministry of Economy and Finance.

The contents of this report are solely the responsibility of the authors and do not necessarily represent the official views of the sponsoring institutions.

1. Introduction

Ischemic stroke is a leading cause of death and disability in both high-income countries and worldwide (1), and the overall disability attributable to stroke (102 million disability-adjusted life-years globally in 2010) is expected to increase over the next few decades (2). Therefore, there is a need to improve the understanding of factors related to the burden of stroke.

It is well-established that initial severity of a stroke episode is an important predictor of the level of disability among stroke survivors (3). Previous studies have described factors that have been associated with initial ischemic stroke severity, including prevalent classical cardiovascular (CV) risk factors, location of the vascular occlusion, size of the thrombus, and stroke etiology, showing that diabetes and hypertension, central occlusions, thrombus >5mm, and cardioembolic etiology predict more severe ischemic strokes (4–7).

A number of studies have demonstrated an association between the physical environment near the home and risk of incident stroke. For example, exposure to ambient fine particulate matter (PM) air pollution (PM_{< 2.5} μm in diameter [PM_{2.5}]) has been associated with an increased risk of hospitalization for ischemic stroke (8–11).

Moreover, environmental noise (especially traffic road noise) has been related to risk of incident cardiovascular events (mainly ischemic heart disease, but also with stroke) (12,13). Conversely, living in neighborhoods with more greenspace (i.e., vegetation) has been linked with lower stroke mortality (14,15).

Interestingly, modifiable characteristics of the social and physical environment have also been associated with initial stroke severity. For example higher socioeconomic

status has been linked to lower stroke severity (16). Some studies have examined the effect of air pollutants and initial stroke severity, but with contradictory results (17,18). We are not aware of any studies considering whether stroke severity is related to either traffic noise or greenspace at the home.

Accordingly, the aim of this study was to assess the association of residential air pollution (PM_{2.5}), road traffic noise and surrounding greenspace with initial stroke severity in patients with acute ischemic stroke. We hypothesized that air pollutants and noise would negatively influence initial ischemic stroke severity whereas exposure to greenspace would be associated with less severe strokes.

2. Methods

The data that support the findings of this study are available from the corresponding author upon reasonable request.

We performed a cross-sectional analysis of the association between residential levels of different environmental factors and initial ischemic stroke severity among patients hospitalized with ischemic stroke. The information regarding environmental and sociodemographic factors was assessed retrospectively at the time of hospital presentation.

2.1 Study design and population

The BASICMAR database (19) is an ongoing prospective register of patients with acute stroke (first episode and recurrence) at University Hospital del Mar, a tertiary public hospital serving an estimated 339,196 residents of two districts (Ciutat Vella and Sant Martí) of the City of Barcelona. Register data are obtained during the index hospital

admission from patients, caregivers, relatives, and/or prior medical records. From this database, we identified 2,786 patients admitted with an episode of acute ischemic stroke (excluding in-hospital strokes, hemorrhagic strokes, and transient ischemic attacks) between January 1, 2005, and December 31, 2014. All patients included in the register were evaluated by a vascular neurologist, with a complete neurovascular examination including imaging studies and diagnostic tests that confirm the diagnosis of acute ischemic stroke. Demographic data and the following vascular risk factors (based on their presence during the index admission, a prior physician diagnosis or need for medical treatment) were recorded from the BASICMAR database: arterial hypertension (evidence of at least two blood pressure measurements $>140/90$ mmHg recorded on different days before stroke onset); diabetes (fasting serum glucose level ≥ 7.0 mmol/L); hyperlipidemia (serum cholesterol levels >220 mg/dL or triglyceride levels >200 mg/dL); atrial fibrillation (AF) confirmed by an ECG performed during admission, previous ischemic heart disease (IHD), defined as previous history of angina pectoris or myocardial infarction; smoking habit; before stroke onset.

2.2 Environmental and socioeconomic data

Patient addresses were geocoded using information from the Cartographic Institute of Catalonia (ICC). Assessment of spatiotemporal exposure was based on a land use regression (LUR) modeling framework developed in the European Study of Cohorts for Air Pollution Effects (ESCAPE) study (20). Following the ESCAPE protocol, measurement sites for $PM_{2.5}$ were selected. These sites included both traffic and background locations, and represented the gradient of various land use, emission sources, and traffic characteristics. Three 2-week monitoring campaigns were conducted in 2009 during different seasons. Estimates were adjusted using data from an ESCAPE

background monitor to account for temporal trends in pollutants between 2009 and the study period (2005-2014) (21,22). GIS (Geographic Information System) data on land uses, traffic indicators, population density, and geographic description of study area were obtained to create potential predictor variables. Multiple linear regression models were constructed following the ESCAPE supervised forward selection protocol using annual average concentrations obtained from the sampling campaign as outcomes. The adjusted R^2 of the final LUR models ranged from 0.71 to 0.85 for the different pollutants, and the cross-validation R^2 ranged from 0.65 to 0.82.

Noise exposure was estimated using Barcelona's municipal strategic noise map (2012) (23). This map follows the recommendations and guidelines of the European Environmental Agency. Exposure to noise (mainly road traffic noise in the study area) was defined as the following four indexes within 50 meters and 250 meters of each home address: L_{day} (from 7 am to 9 pm), $L_{evening}$ (from 9 pm to 11 pm), L_{night} (from 11 pm to 7 am) and L_{den} (Day-evening-night-weighted). The L_{day} , $L_{evening}$ and L_{night} indexes are A-weighted sound levels, determined over the corresponding day, evening and night periods. L_{den} indicator is an average sound pressure level over all days, evenings and nights in a year. Indexes values (dB) at the street nearest to each participant's residential address were assigned to that participant.

To characterize residential surrounding greenspace, we used the Normalized Difference Vegetation Index (NDVI) derived from the Landsat 8 Operational Land Imager (OLI) sensor data at 30m x 30m resolution. Given that the maximum vegetation in our study region occurs in the spring and autumn months, the Landsat image from April 16, 2013 was selected in order to maximize the contrast in exposure (24). The findings of our previous studies support the stability of the NDVI spatial contrast over years in the

study region (24). Usually, areas of barren rock, sand, or snow usually show very low NDVI values (<0.1) and sparse vegetation such as grasslands result in moderate NDVI values (0.2 to 0.5). Residential surrounding greenspace was quantified as the average NDVI within 100, 300, and 500 meters around each participant's residential geocoded address.

Figure 1 represents a map of the city of Barcelona where intensity of brown color depicts PM_{2.5} concentration (corresponding to the standard road map) and the intensity of green color, NDVI concentration.




Figure 1. Map of the city of Barcelona. The red square outlines the two districts assessed. Blue dots

represent patients. Intensity of brown color depicts PM_{2.5} concentration (corresponding to the standard road map) and the intensity of green color, NDVI concentration.

Urban Vulnerability Index is a measure of neighborhood socioeconomic status (NSS) at the census tract level (median area of 0.08 km² for the study area) based on 21 indicators of urban vulnerability grouped into four themes: sociodemographic vulnerability (five indicators), socioeconomic vulnerability (six indicators), housing vulnerability (five indicators), and subjective perception of vulnerability (five indicators) (25). This measure of NSS has been widely used in previous studies and has showed strong correlations with individual socioeconomic status indicators such as “temporary employment”, “unskilled work” and “illiteracy” (26,27).

2.3 Stroke severity outcome

The primary outcome for this analysis was initial stroke severity measured by the National Institute of Health Stroke Scale (NIHSS; 0-42) score, considering the higher the score, the greater the stroke severity. The NIHSS is a 15-item neurologic examination stroke scale used to evaluate the effect of stroke on the levels of consciousness, language, neglect, visual-field loss, extraocular movement, motor strength, ataxia, dysarthria, and sensory loss. A trained observer rates the patient’s ability to answer questions and perform activities. Ratings for each item are scored with 3 to 5 grades with 0 as normal, and there is an allowance for untestable items (see Supplementary material). Moderate to severe stroke was defined as NIHSS score >5 based on previous studies (3,28,29). The NIHSS score was assessed by the neurologist in charge once the patient arrived at the emergency room and every score was recorded from the BASICMAR database.

2.4 Statistical analysis

We performed a descriptive analysis of the sociodemographic and clinical variables calculating percentages for categorical variables and mean with corresponding standard deviations (or median with interquartile range in case of not normally distributed) for quantitative variables.

We considered all the environmental exposure variables (air pollutant concentration $PM_{2.5}$, every noise indicator- *Lday*, *Levening*, *Lnight* and *Lden*- and average NDVI for 100, 300 and 500 meters) modeled as quartiles for the main analysis. Analysis considering all the exposure variables modeled as linear continuous (interquartile range increments) were also performed.

We used logistic and negative binomial regression models to evaluate the association between $PM_{2.5}$, traffic noise, greenspace and severity (considered either as dichotomously NIHSS score >5 and as continuous variable). We modeled each association for each environmental factor as follows: a) Model 1, adjusted for age, sex, smoking, and NSS; b) Model 2, Model 1 plus cardiovascular risk factors and previous history of coronary heart disease or stroke; c) Model 3, a multi-environmental model ($PM_{2.5}$, road traffic noise and greenspace), adjusted for the rest of covariables (age, sex, smoking, NSS, cardiovascular risk factors, previous history of coronary heart disease or stroke). Noise indicators were considered individually in each model (i.ea separate model for each noise indicator). Models were not adjusted for stroke etiology as etiology is potentially a causal mediator between environmental factors and stroke severity.

Regarding greenspace models, the main analysis was done using NDVI buffer within 300 meters; sensitivity analyses were also performed using NDVI buffer within 100 and 500 meters.

2.5 Ethical consideration

The information used in this study was collected from the prospective BASICMAR register, with the approval of our local ethics committee. All patients (or their proxies) provided informed consent prior to their inclusion in the study registry.

3. Results

3.1 Clinical and environmental characteristics

Of the 2,786 patients hospitalized with acute ischemic stroke from the Ciutat Vella and Sant Martí district and included in the BASICMAR register during the study period, 25 were excluded due to admission date errors or unknown (9 patients) or missing initial NIHSS (16 patients). The final sample included 2,761 patients with ischemic stroke.

Sociodemographic, clinical and environmental characteristics are described in Table 1.

Sociodemographic characteristics	Mean (SD) or Median [IQR] or n (%)
Age, y	75 (12.5)
Female	1399 (50.7)
Neighborhood Socioeconomic Status*	0.68 (0.20)
Cardiovascular risk factors and comorbidities	
Current smoker	568 (20.6)
Hypertension	2111 (76.5)
Diabetes Mellitus	958 (34.7)
Dyslipidemia	1239 (44.9)
Previous history of coronary heart disease	429 (15.5)
Previous history of stroke	550 (20)
Atrial Fibrillation	719 (26)
Outcome	
NIHSS	5 [3-12]
NIHSS >5	1267 (45.9)
Environmental characteristics	Mean (SD) or Median [IQR]
PM _{2.5} (µg/m ³)	17.89 [15.20-21.88]

NDVI average**	
100 m	0.154 [0.125-0.199]
300 m	0.166 [0.136-0.204]
500 m	0.170 [0.138-0.200]
Road traffic noise level at nearest street (dB)	
<i>LDay</i>	63 (4.9)
<i>LEvening</i>	61 (4.9)
<i>LNight</i>	55 (5.5)
<i>Lden</i>	65 (5)

Table 1. Sociodemographic characteristics of patients with acute ischemic stroke (n=2761)

* NSS index where 1 more deprived and 0 less deprived. **NDVI average within 100, 300 and 500 meters buffer where 1 is most greener and 0, less

The majority of the patients lived in moderately disadvantaged neighborhoods.

Hypertension was the most frequent cardiovascular risk factor (76.5%). There was only a high correlation across noise indicators, not between any other exposures metric (Table 2).

	PM _{2.5}	NDVI300	<i>Lday</i>	<i>Levening</i>	<i>Lnight</i>
PM _{2.5}	1				
NDVI300	-0.12	1			
<i>Lday</i>	0.24	-0.08	1		
<i>Levening</i>	0.24	-0.18	0.94	1	
<i>Lnight</i>	0.25	-0.15	0.92	0.94	1
<i>Lden</i>	0.25	-0.13	0.97	0.96	0.98

Table 2 Spearman correlation matrix between all pairs of the exposures. All correlations were statistically significant. PM_{2.5}: particulate matter 2.5 microns; NDVI 300: Normalized Difference Vegetation Index within 300 meters; *Lday*:day noise indicator; *Levening*: evening noise indicator; *Lnight*: night noise indicator; *Lden*: average noise indicator.

3.2 Association of environmental factors on initial stroke severity

Higher residential surrounding greenspace (4th quartile compared to 1st quartile) in 300m buffer was associated with less severe acute ischemic stroke (OR of NIHSS>5, 0.75; 95% CI 0.60-0.95). Conversely, living in areas with higher average noise (*Lden*, 4th quartile compared to 1st quartile) had a higher risk of severe stroke (OR, 1.30; 95% CI: 1.02-1.65). Residential PM_{2.5} was not associated with initial stroke severity (see Table 3 and 4). When adjusting for PM_{2.5} levels and *Lden*, the protective effect of greenspace persisted (Table5).

Exposure	Stroke severity (NIHSS >5)	
	OR (95% CI)	
	Model 1	Model 2
Annual PM _{2.5}		
Q2	1.03 (0.82-1.30)	1.01 (0.80-1.26)
Q3	1.00 (0.80-1.23)	0.93 (0.74-1.17)
Q4	1.10 (0.88-1.40)	1.04 (0.83-1.31)
<i>LDay</i>		
Q2	1.11 (0.90-1.40)	1.10 (0.88-1.37)
Q3	1.10 (0.90-1.35)	1.10 (0.88-1.35)
Q4	1.31 (1.01-1.70)	1.30 (0.99-1.67)
<i>LEvening</i>		
Q2	1.02 (0.82-1.30)	1.00 (0.80-1.36)
Q3	1.16 (0.93-1.44)	1.14 (0.91-1.42)
Q4	1.30 (1.04-1.63)	1.27 (1.01-1.60)
<i>LNight</i>		
Q2	1.04 (0.83-1.30)	1.04 (0.83-1.31)
Q3	1.14 (0.91-1.41)	1.12 (0.90-1.4)
Q4	1.23 (1.00-1.54)	1.23 (0.98-1.55)
<i>Lden</i>		
Q2	1.05 (0.85-1.30)	1.05 (0.85-1.3)
Q3	1.15 (0.91-1.46)	1.15 (0.90-1.46)
Q4	1.30 (1.03-1.65)	1.30 (1.02-1.65)
NDVI 300 meters		
Q2	1.01 (0.80-1.26)	1.03 (0.81-1.30)

Q3	0.70 (0.56-0.88)	0.71 (0.60-0.90)
Q4	0.76 (0.60-0.95)	0.75 (0.60-0.95)

Table 3. Odds ratio (OR) and 95% confidence interval of being hospitalized with a severe ischemic stroke defined as an initial NIH Stroke Scale score of >5. PM_{2.5}, noise indicators and residential surrounding greenspace at 300 meters, all modelled as quartiles variables, with Q1 being the lowest concentration/value and taken as reference. Stroke severity modeled as dichotomous variable NIHSS >5. Model 1: adjusted by age sex, smoking status and NSS; Model 2: model 1 plus hypertension, diabetes mellitus, dyslipidemia, prior history of coronary heart disease, prior history of stroke/transient ischemic attack. Test of linear trend across quartiles: PM_{2.5} p= 0.469; L_{Day} p=0.053 L_{Evening} p= 0.016 L_{Night} p=0.068 L_{den} p=0.027 NDVI₃₀₀ p= 0.010

Exposure	Stroke severity (NIHSS) B Coefficient (95% CI)	
	Model 1	Model 2
Air pollutant concentration		
Annual PM _{2.5}		
Q2	-0.14 (-0.11-0.08)	-0.01(-0.10-0.08)
Q3	0.02 (-0.07-0.11)	0.00(-0.09-0.97)
Q4	0.12 (-0.08-0.11)	0.00(-0.09-0.09)
Road traffic noise		
L _{Day}		
Q2	0.066 (-0.02-0.16)	0.06 (-0.02-0.15)
Q3	0.09 (0.00-0.18)	0.09 (0.00-0.18)
Q4	0.09(-0.016-0.20)	0.08 (-0.02-0.19)
L _{Evening}		
Q2	0.03 (-0.06-0.13)	0.03 (-0.06-0.13)
Q3	0.09 (0.00-0.18)	0.09 (-0.00-0.18)
Q4	0.08 (-0.01-0.18)	0.07 (-0.01-0.17)
L _{Night}		
Q2	0.04 (-0.05-0.13)	0.04 (-0.05-0.13)
Q3	0.08 (-0.01-0.17)	0.07 (-0.02-0.16)
Q4	0.08 (-0.01 (0.17)	0.08 (-0.01-0.17)
L _{den}		
Q2	0.05 (-0.04-0.14)	0.05 (-0.04 -0.14)

Q3	0.09 (-0.01-0.19)	0.08 (-0.01-0.18)
Q4	0.09 (-0.00-0.19)	0.09 (-0.01-0.19)
Residential surrounding greenspace		
NDVI 300 meters		
Q2	0.01 (-0.08-0.10)	-0.01 (-0.08-0.11)
Q3	-0.08 (-0.18-0.01)	-0.08 (-0.17-0.015)
Q4	-0.09 (-0.19- -0.00)	-0.11 (-0.21- - 0.01)

Table 4. Association between environmental factors and initial NIH Stroke Scale score as a continuous variable using negative binomial regression. PM_{2.5}, noise indicators and residential surrounding greenspace at 300 meters, all modelled as quartiles variables, with Q1 being the lowest concentration/value and taken as reference. Stroke severity modeled as continuous. Model 1: adjusted by age sex, smoking status and NSS; Model 2: model 1 plus hypertension, diabetes mellitus, dyslipidemia, prior history of coronary heart disease, prior history of stroke/transient ischemic attack. Test of linear trend across quartiles: PM2.5 p= 0.286; *LDay* p=0.018 *LEvening* p= 0.017 *LNight* p=0.069 *Lden* p=0.025 NDVI_300 p= 0.015

Exposure	Model 3	
	Dichotomous NIHSS >5	Change in NIHSS
	OR (95% CI)	(95% CI)
Air pollutant concentration		
Annual PM _{2.5}		
Q2	0.97 (0.77-1.21)	-0.03 (-0.13-0.06)
Q3	0.88 (0.7-1.11)	-0.02 (-0.11-0.07)
Q4	0.95 (0.75-1.20)	-0.03 (-0.13-0.06)
Road traffic noise		
<i>Lden</i>		
Q2	1.04 (0.84-1.30)	0.05 (-0.04-0.14)
Q3	1.16 (0.91-1.48)	0.09 (-0.01-0.19)
Q4	1.19 (0.93-1.53)	0.06 (-0.13-0.06)
Residential surrounding greenspace		
NDVI 300 meters		
Q2	1.00 (0.79-1.27)	0.01 (-0.09-0.10)
Q3	0.71 (0.56-0.90)	-0.08 (-0.18-0.01)
Q4	0.76 (0.60-0.95)	-0.11 (-0.21-0.014)

Table 5. Results of logistic and binomial regression multienvironmental model of the association between exposure to environmental factors and stroke severity. PM_{2.5}, *Lden* indicator and residential surrounding greenspace at 300 meters, all modelled as quartiles variables, with Q1 being the lowest concentration/value and taken as reference. Stroke severity modeled either as dichotomous NIHSS >5 as continuous. Model 3: model 2 plus all environmental exposures (PM_{2.5}, *Lden* and NDVI 300 meters)

Sensitivity analyses considering NDVI within 500 meters buffer were consistent with those of the main analysis. Results considering NDVI within a 100 meters buffer were in the same direction, but not statistically significant (see Appendices Table A1 and Table A2a and A2b). Results considering environmental exposure variables modeled as linear continuous variables (IQR increments) were similar to the main analysis (see Appendices Table A3 and Table A4a and A4b).

4. Discussion

To our knowledge, this is the first study simultaneously evaluating the influence of an array of urban-related exposures on initial stroke severity which also adds to the scarce available evidence on environmental determinants of this health endpoint. The individuals studied were part of a clinically well-described sample of patients with acute ischemic stroke, which facilitated to adjust for different factors (age, CV risk factors and NSS) that have been previously described to be related with severity. We observed that higher residential surrounding greenspace had a potential beneficial influence on initial stroke severity. This association persisted after adjusting for other environmental factors (PM_{2.5} and noise) in multi-exposure models. Exposure to noise was associated with more severe initial stroke, however, the association did not persist after adjustment for other environmental factors (PM_{2.5} and greenspace). For air pollution our findings were not conclusive.

There is growing evidence supporting the beneficial health influences exposure to greenspace. This exposure has been associated with improved mental and physical health, including lower risk of mortality and morbidity (30). For instance, higher residential surrounding greenspace has been associated with lower risk of

cardiovascular mortality (31) and clinical visits for cardiovascular problems (32); however, there are inconsistencies in these reported associations (33).

Residential proximity greenspace has been shown to be related with lower stroke admissions (34) and with higher survival rates after acute ischemic stroke (14,15). However, the available evidence on the association between this exposure and stroke is still very scarce, with no study reporting on the association with severity of stroke.

Different mechanisms have been hypothesized for how exposure to greenspace might have a beneficial effect on health. On one hand, it might promote stress reduction (35), social interactions (36) and physical activity (32). In fact, a recent study found an association between prestroke regular physical activity and initial stroke severity (37). On the other, exposure to greenspace has been associated with an immunoregulatory effect (38), lower levels of sympathetic activation, reduce oxidative stress, and higher angiogenic capacity (39). These mechanisms could potentially be implicated in the initial formation of the thrombus and the systemic inflammatory response to stroke, and thus, initial severity. However, there is a need for further research in this area to understand the possible mechanisms by which prestroke exposure to greenspace would influence stroke severity.

In this study we also observed that patients living in areas with higher annual average noise values had a 20% higher risk of severe stroke, compare to those living in areas with lower average noise values. This association did not persist after adjustment for NDVI and air pollution.

Road traffic noise has also been related with adverse CV outcomes (13), but few studies have assessed the influence on stroke, suggesting that traffic noise might play a

deleterious role increasing its risk (40). There are no studies, though, testing specifically its effect in acute ischemic stroke severity. A possible mechanism that could explain the association is that noise may also lead to endothelial dysfunction and arterial hypertension (41).

Air pollution has been related with higher risk of acute ischemic stroke events (42,43) and higher mortality (10). We did not find any association between exposure to long-term air pollutant concentrations ($PM_{2.5}$) and stroke severity. There are few studies that have tested the effect on severity and results are contradictory (18,44). Our results were in line with a previously study performed in a similar cohort in another urban setting (London) that showed no association between air pollution exposure and stroke severity (45).

Our study had several limitations. The urban environment across the two districts of Barcelona where patients lived offered limited variability in exposure to environmental factors (Figure 1). On the other hand, despite the different timeframes from which we obtained the data from noise map indicators, satellite-based NDVI values and spatial estimates of air pollutant levels generated by land-use regression models, we effectively assumed that the city spatial surface and the spatial distribution of (road traffic) noise, greenspaces and air pollutants remained constant over the study period. There are some reports supporting the stability of the spatial contrasts for noise (46), greenspace (47,48) and air pollution (21,22) in the study region. Furthermore, to our knowledge, there was no major change in land use, emissions profiles, or traffic flow between the year of land-use regression model construction (2009) and the years of our study (2005-2014).

Our use of satellite-based NDVI to assess greenspace surrounding participants' homes enabled us to take account of small-area green spaces (e.g. roadside trees and home gardens) in a standardized way; however, NDVI does not distinguish between different types of vegetation (e.g. trees, shrubs, grasses, etc.) which could have been relevant in our evaluated association. NDVI measures do not take the quality or frequency of use of green spaces into account. Quality characteristics of green spaces like aesthetics, biodiversity, walkability, sport/play facilities, safety, and organized social events have been suggested to predict the use of green spaces (49) and could have an effect on our investigated associations.

We did not have data regarding individual socioeconomic status. This factor might have influence in patients' lifestyle behavior (and, for example, individuals with higher socioeconomic status might be prone to live nearby more dense green areas). However, the adjustment for a robust index of area vulnerability (including 21 items) and a set of comorbidities strongly related to social class might have reduced the likelihood for a potential residual confounding for social class.

We have considered as initial severity the first NIHSS score assessed once the patient arrived at the emergency room. It is known that ischemic stroke is an acute disease and its severity is mainly established at the very beginning of the episode and driven by the location and characteristic of the arterial thrombus. However, it is possible that some stroke patients could initially present with fluctuating symptoms and that other factors such as distance to the hospital, living conditions, or the presence of a bystander could have influence the initial score assessment of the stroke severity.

5. Conclusion

Our study found that, in an urban setting, residential surrounding greenspace and traffic noise were associated with initial stroke severity, suggesting an important influence of the built environment on the global burden of stroke. Further studies are needed to assess which aspects of exposure to neighborhood green spaces could influence the initial severity of ischemic stroke.

Disclosures: None

References

1. Global, regional, and national burden of neurological disorders during 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Neurol.* England; 2017 Nov;16(11):877–97.
2. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Bennett DA, Moran AE, et al. from the Global Burden of Disease Study 2010. 2014;383(9913):245–54.
3. Adams HPJ, Davis PH, Leira EC, Chang KC, Bendixen BH, Clarke WR, et al. Baseline NIH Stroke Scale score strongly predicts outcome after stroke: A report of the Trial of Org 10172 in Acute Stroke Treatment (TOAST). *Neurology.* United States; 1999 Jul;53(1):126–31.
4. Cuadrado-Godia E, Jimena S, Ois A, Rodriguez-Campello A, Giralt-Steinhauer E, Soriano-Tarraga C, et al. Factors associated with early outcome in patients with large-vessel carotid strokes. *J Neurol Neurosurg Psychiatry.* England; 2013 Mar;84(3):305–9.

5. Fischer U, Arnold M, Nedeltchev K, Brekenfeld C, Ballinari P, Remonda L, et al. NIHSS score and arteriographic findings in acute ischemic stroke. *Stroke*. United States; 2005 Oct;36(10):2121–5.
6. Hong K-S, Lee J, Bae H-J, Lee JS, Kang D-W, Yu K-H, et al. Greater stroke severity predominates over all other factors for the worse outcome of cardioembolic stroke. *J Stroke Cerebrovasc Dis*. United States; 2013 Nov;22(8):e373-80.
7. Marder VJ, Chute DJ, Starkman S, Abolian AM, Kidwell C, Liebeskind D, et al. Analysis of thrombi retrieved from cerebral arteries of patients with acute ischemic stroke. *Stroke*. United States; 2006 Aug;37(8):2086–93.
8. Wellenius GA, Schwartz J, Mittleman MA. Air pollution and hospital admissions for ischemic and hemorrhagic stroke among medicare beneficiaries. *Stroke*. 2005;36(12):2549–53.
9. Wellenius GA, Burger MR, Coull BA, Schwartz J, Suh HH, Koutrakis P, et al. Ambient air pollution and the risk of acute ischemic stroke. *Arch Intern Med* [Internet]. 2012;172(3):229–34. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22332153>
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3639313>
10. Scheers H, Jacobs L, Casas L, Nemery B, Nawrot TS. Long-Term Exposure to Particulate Matter Air Pollution Is a Risk Factor for Stroke. *Stroke* [Internet]. 2015 Oct 26;46(11):3058 LP-3066. Available from: <http://stroke.ahajournals.org/content/46/11/3058.abstract>

11. Huang F, Luo Y, Guo Y, Tao L, Xu Q, Wang C, et al. Particulate Matter and Hospital Admissions for Stroke in Beijing, China: Modification Effects by Ambient Temperature. *J Am Heart Assoc* [Internet]. 2016;5(7):1–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27413045>
12. Kempen E van, Casas M, Pershagen G, Foraster M. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Cardiovascular and Metabolic Effects: A Summary. *Int J Environ Res Public Health*. Switzerland; 2018 Feb;15(2).
13. Cai Y, Hodgson S, Blangiardo M, Gulliver J, Morley D, Fecht D, et al. Road traffic noise, air pollution and incident cardiovascular disease: A joint analysis of the HUNT, EPIC-Oxford and UK Biobank cohorts. *Environ Int*. Netherlands; 2018 May;114:191–201.
14. Wilker EH, Wu C-D, McNeely E, Mostofsky E, Spengler J, Wellenius GA, et al. Green space and mortality following ischemic stroke. *Environ Res*. Netherlands; 2014 Aug;133:42–8.
15. Vienneau D, de Hoogh K, Faeh D, Kaufmann M, Wunderli JM, Rösli M. More than clean air and tranquillity: Residential green is independently associated with decreasing mortality. *Environ Int*. Elsevier; 2017;108(July):176–84.
16. Rey V, Faouzi M, Huchmand-Zadeh M, Michel P. Stroke initial severity and outcome relative to insurance status in a universal health care system in Switzerland. *Eur J Neurol*. England; 2011 Aug;18(8):1094–7.
17. Maheswaran R, Pearson T, Beevers SD, Campbell MJ, Wolfe CD. Air Pollution

- and Subtypes, Severity and Vulnerability to Ischemic Stroke-A Population Based Case-Crossover Study. *PLoS One*. United States; 2016;11(6):e0158556.
18. Wing JJ, Sanchez BN, Adar SD, Meurer WJ, Morgenstern LB, Smith MA, et al. Synergism of Short-Term Air Pollution Exposures and Neighborhood Disadvantage on Initial Stroke Severity. *Stroke*. United States; 2017 Nov;48(11):3126–9.
 19. Roquer J, Rodríguez-Campello A, Gomis M, Jiménez-Conde J, Cuadrado-Godia E, Vivanco R, et al. Acute stroke unit care and early neurological deterioration in ischemic stroke. *J Neurol* [Internet]. 2008;255(7):1012–7. Available from: <http://dx.doi.org/10.1007/s00415-008-0820-z>
 20. Beelen R, Raaschou-Nielsen O, Stafoggia M, Andersen ZJ, Weinmayr G, Hoffmann B, et al. Effects of long-term exposure to air pollution on natural-cause mortality: An analysis of 22 European cohorts within the multicentre ESCAPE project. *Lancet*. 2014;383(9919):785–95.
 21. Cesaroni G, Porta D, Badaloni C, Stafoggia M, Eeftens M, Meliefste K, et al. Nitrogen dioxide levels estimated from land use regression models several years apart and association with mortality in a large cohort study. *Environ Health*. England; 2012 Jul;11:48.
 22. Eeftens M, Beelen R, Fischer P, Brunekreef B, Meliefste K, Hoek G. Stability of measured and modelled spatial contrasts in NO₂ over time. *Occup Environ Med*. England; 2011 Oct;68(10):765–70.
 23. Tel B. Resum mapes estratègics de soroll (2012-2017). 2017;2017:523–5.

24. Dadvand P, Sunyer J, Basagana X, Ballester F, Lertxundi A, Fernandez-Somoano A, et al. Surrounding greenness and pregnancy outcomes in four Spanish birth cohorts. *Environ Health Perspect.* United States; 2012 Oct;120(10):1481–7.
25. Atlas of urban vulnerability, Ministry of Public Works and Infrastructure, Spain. 2012
https://www.fomento.gob.es/MFOM/LANG_CASTELLANO/DIRECCIONES_GENERALES/ARQ_VIVIENDA/SUELO_Y_POLITICAS/OBSERVATORIO/Atlas_Vulnerabilidad_Urbana.
26. Bosch de Basea M, Figuerola J, Pardina M, Espinosa A, Cardis E, Vilar J, et al. CT scan exposure in Spanish children and young adults by socioeconomic status: Cross-sectional analysis of cohort data. *PLoS One.* 2018;13(5):e0196449.
27. Valentin A, Maitre L, McEachan R, Lepeule J, Ambros A, Nieuwenhuijsen M, et al. The Urban Exposome during Pregnancy and Its Socioeconomic Determinants. *Environ Health Perspect.* 2018;126(7):77005.
28. Khatri P, Tayama D, Cohen G, Lindley RI, Wardlaw JM, Yeatts SD, et al. Effect of intravenous recombinant tissue-type plasminogen activator in patients with mild stroke in the third international stroke trial-3: Post hoc analysis. *Stroke.* 2015;46(8):2325–7.
29. Sandercock P, Wardlaw JM, Lindley RI, Dennis M, Cohen G, Murray G, et al. The benefits and harms of intravenous thrombolysis with recombinant tissue plasminogen activator within 6 h of acute ischaemic stroke (the third international stroke trial [IST-3]): A randomised controlled trial. *Lancet* [Internet]. Elsevier Ltd; 2012;379(9834):2352–63. Available from:

[http://dx.doi.org/10.1016/S0140-6736\(12\)60768-5](http://dx.doi.org/10.1016/S0140-6736(12)60768-5)

30. Kondo MC, Fluehr JM, McKeon T, Branas CC. Urban green space and its impact on human health. *Int J Environ Res Public Health*. 2018;15(3).
31. Gascon M, Triguero-Mas M, Martinez D, Dadvand P, Rojas-Rueda D, Plasencia A, et al. Residential green spaces and mortality: A systematic review. *Environ Int*. Netherlands; 2016 Jan;86:60–7.
32. Maas J, Verheij RA, Spreeuwenberg P, Groenewegen PP. Physical activity as a possible mechanism behind the relationship between green space and health: a multilevel analysis. *BMC Public Health*. England; 2008 Jun;8:206.
33. Tamosiunas A, Grazuleviciene R, Luksiene D, Dedele A, Reklaitiene R, Baceviciene M, et al. Accessibility and use of urban green spaces, and cardiovascular health: findings from a Kaunas cohort study. *Environ Health*. England; 2014 Mar;13(1):20.
34. Foster S, Giles-Corti B, Knuijman M, Martin K, Christian H, Boruff BJ, et al. The association between neighborhood greenness and cardiovascular disease: an observational study. *BMC Public Health* [Internet]. ???; 2012;12(1):1. Available from: ???
35. Triguero-Mas M, Donaire-Gonzalez D, Seto E, Valentin A, Martinez D, Smith G, et al. Natural outdoor environments and mental health: Stress as a possible mechanism. *Environ Res*. Netherlands; 2017 Nov;159:629–38.
36. Maas J, van Dillen SME, Verheij RA, Groenewegen PP. Social contacts as a possible mechanism behind the relation between green space and health. *Health*

Place. England; 2009 Jun;15(2):586–95.

37. Reinholdsson M, Palstam A, Sunnerhagen KS. Prestroke physical activity could influence acute stroke severity (part of PAPSIGOT). *Neurology* [Internet]. 2018;0:10.1212/WNL.0000000000006354. Available from: <http://www.neurology.org/lookup/doi/10.1212/WNL.0000000000006354>
38. Rook GA. Regulation of the immune system by biodiversity from the natural environment: An ecosystem service essential to health. *Proc Natl Acad Sci* [Internet]. 2013;110(46):18360–7. Available from: <http://www.pnas.org/cgi/doi/10.1073/pnas.1313731110>
39. Yeager R, Riggs DW, DeJarnett N, Tollerud DJ, Wilson J, Conklin DJ, et al. Association Between Residential Greenness and Cardiovascular Disease Risk. *J Am Heart Assoc* [Internet]. 2018;7(24). Available from: <https://www.ahajournals.org/doi/10.1161/JAHA.118.009117>
40. Halonen JI, Hansell AL, Gulliver J, Morley D, Blangiardo M, Fecht D, et al. Road traffic noise is associated with increased cardiovascular morbidity and mortality and all-cause mortality in London. *Eur Heart J*. 2015;36(39):2653–61.
41. Munzel T, Schmidt FP, Steven S, Herzog J, Daiber A, Sorensen M. Environmental Noise and the Cardiovascular System. *J Am Coll Cardiol*. United States; 2018 Feb;71(6):688–97.
42. Wang Y, Eliot MN, Wellenius GA. Short-term changes in ambient particulate matter and risk of stroke: A systematic review and meta-analysis. *J Am Heart Assoc*. 2014;3(4):1–23.

43. Vivanco-Hidalgo RM, Wellenius G, Basagaña X, Cirach M, González AG, Ceballos P de, et al. Short-term exposure to traffic-related air pollution and ischemic stroke onset in Barcelona, Spain. *Environ Res [Internet]*. Elsevier Inc.; 2018;162(December 2017):160–5. Available from: <https://doi.org/10.1016/j.envres.2017.12.024>
44. Andersen ZJ, Olsen TS, Andersen KK, Loft S, Ketzel M, Raaschou-Nielsen O. Association between short-term exposure to ultrafine particles and hospital admissions for stroke in Copenhagen, Denmark. *Eur Heart J*. 2010;31(16):2034–40.
45. Maheswaran R, Pearson T, Beevers SD, Campbell MJ, Wolfe CD. Air pollution and subtypes, severity and vulnerability to ischemic stroke—a population based case-crossover study. *PLoS One*. 2016;11(6).
46. Dadvand P, Ostro B, Figueras F, Foraster M, Basagaña X, Valentín A, et al. Residential proximity to major roads and term low birth weight: The roles of air pollution, heat, noise, and road-adjacent trees. *Epidemiology*. 2014;25(4):518–25.
47. Nieuwenhuijsen MJ, Gascon M, Martinez D, Ponjoan A, Blanch J, Garcia-Gil MDM, et al. Air Pollution, Noise, Blue Space, and Green Space and Premature Mortality in Barcelona: A Mega Cohort. *Int J Environ Res Public Health*. Switzerland; 2018 Oct;15(11).
48. Dadvand P, Wright J, Martinez D, Basagana X, McEachan RRC, Cirach M, et al. Inequality, green spaces, and pregnant women: roles of ethnicity and individual and neighbourhood socioeconomic status. *Environ Int*. Netherlands; 2014

Oct;71:101–8.

49. McCormack GR, Rock M, Toohey AM, Hignell D. Characteristics of urban parks associated with park use and physical activity: a review of qualitative research. *Health Place. England*; 2010 Jul;16(4):712–26.