

EAACI Biologicals Guidelines – Omalizumab for the treatment of chronic spontaneous urticaria in children and in adults

Authors: Ioana Agache¹, Cezmi Akdis^{2,3}, Mubeccel Akdis², Knut Brockow⁴, Tomas Chivato⁵, Stefano del Giacco⁶, Thomas Eiwegger^{7,8,9}, Kilian Eyerich¹⁰, Ana Giménez-Arnau¹¹, Jan Gutermuth¹², Emma Guttman-Yassky¹³, Marcus Maurer¹⁴, Graham Ogg¹⁵, Peck Y. Ong¹⁶, Liam O'Mahony¹⁷, Jürgen Schwarze¹⁸, Amena Warner¹⁹, Thomas Werfel²⁰, Oscar Palomares^{21*}, Marek Jutel^{22,23*}

* Joint last authorship

External peer review: **please suggest**

Corresponding author: Ioana Agache; 2A, Pictor Ion Andreescu, Brasov, Romania, 500051

Affiliations

¹Transylvania University, Faculty of Medicine, Brasov, Romania

² Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland

³ Christine-Kühne-Center for Allergy Research and Education (CK-CARE), Davos, Switzerland

⁴ Department of Dermatology and Allergology Biederstein, School of Medicine, Technical University of Munich, Munich, Germany

⁵ School of Medicine University CEU San Pablo, Madrid, Spain

⁶ Department of Medical Sciences and Public Health, University of Cagliari, Italy

⁷ Translational Medicine Program, Research Institute, Hospital for Sick Children, Toronto, ON, Canada

⁸ Department of Immunology, University of Toronto, Toronto, ON, Canada

⁹ Division of Immunology and Allergy, Food Allergy and Anaphylaxis Program, The Hospital for Sick Children, Departments of Paediatrics and Immunology, University of Toronto, Toronto, Canada

¹⁰ Department of Dermatology and Allergy Biederstein, Technical University of Munich, Munich, Germany

¹¹ Department of Dermatology, Hospital del Mar- Institut Mar d'Investigacions Mèdiques, Universitat Autònoma de Barcelona, Spain

¹² Department of Dermatology, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel (VUB), Brussels, Belgium

¹³ Department of Dermatology, Icahn School of Medicine at Mount Sinai, New York, USA

¹⁴ Dermatological Allergology, Allergie-Centrum-Charité, Department of Dermatology and Allergy, Charité - Universitätsmedizin Berlin, Germany

¹⁵ MRC Human Immunology Unit, MRC Weatherall Institute of Molecular Medicine, Oxford NIHR Biomedical Research Centre, Radcliffe Department of Medicine, University of Oxford, UK

¹⁶ Division of Clinical Immunology & Allergy, Children's Hospital Los Angeles, Keck School of Medicine, University of Southern California, USA

¹⁷ Departments of Medicine and Microbiology, APC Microbiome Ireland, University College Cork, Cork, Ireland

¹⁸ Centre for Inflammation Research, Child Life and Health, The University of Edinburgh, Edinburgh, United Kingdom

¹⁹ Allergy UK

²⁰ Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany

²¹ Department of Biochemistry and Molecular Biology, Chemistry School, Complutense University of Madrid, Spain

²² University of Wrocław, Department of Clinical Immunology, Wrocław, Poland

Short title: EAACI Biologicals Guidelines – omalizumab for chronic spontaneous urticaria

Acknowledgements: The GDG is grateful to all the methodology team from the Iberoamerican Cochrane Center (Biomedical Research Institute Sant Pau) who conducted the systematic reviews for efficacy, safety and economic evidence.

Abstract (unstructured, 200 words)

Chronic spontaneous urticaria (CSU) imposes a significant burden on patients, families and healthcare systems. Management is difficult, due to disease heterogeneity and complexity in care pathways. Better understanding of the mechanisms has enabled a stratified approach to the management of CSU, supporting the use of targeted treatment with omalizumab. However, there are still many issues that require further clarification. These include selection of responders, the definition of response, strategies to enhance the responder rate, the duration of treatment and its regimen (in the clinic or home-based) and its cost-effectiveness. The EAACI Guidelines on the use of biologicals in CSU follow the GRADE approach in formulating recommendations for each biological and each outcome. In addition, future approaches and research priorities are discussed.

Key words: chronic spontaneous urticaria, GRADE, guidelines, omalizumab

Abbreviations

AE = adverse events
AIS = activity impairment score
ASST = autologous serum skin test
BHRA = basophil histamine release assay
BTs = basophil tests
CI = confidence interval
CIU = chronic idiopathic urticaria
CSU = chronic spontaneous urticaria
CU-Q2oL = Chronic Urticaria Quality of Life Questionnaire
DARPs = Designed Ankyrin Repeat Proteins
DLQI = Dermatology Life Quality Index
EAACI = European Academy of Allergy and Clinical Immunology
ECs = Endothelial cells
EMA = European Medicine Agency
EtD = evidence to decision
FcεR1 = high-affinity IgE receptor
FDA = Food and Drug administration
FcεRI = high affinity receptors for IgE
GDG = Guideline Development Group
GPCRs = G protein-coupled receptors
GRADE = Grading of Recommendations Assessment, Development and Evaluation
HCP = healthcare professionals
Ig = immunoglobulin
IL = interleukin
ISS = itch severity score
IRR = incidence rate ratios
ITAMs = immunoreceptor tyrosine-based activation motifs
mAb = monoclonal antibodies

MCs = mast cells
MD = mean difference
MID = minimal important difference
MMP= metalloproteinase
MRGPRX2 = Mas-related G protein-coupled receptor X2
QoL = quality of life
PICO = population, intervention, comparator, and outcomes
RCT = randomised controlled trial
ROB = risk of bias
RR= risk ratio
SAE = serious adverse events
SC= subcutaneous
SOF = summary of findings
SR = systematic review
SYK = spleen tyrosine kinase
TSLP = thymic stromal lymphopoietin
UAS = urticaria activity score
UCT = urticaria control score
WIS = work impairment score
WPAI = work productivity and activity impairment

I. Introduction

a. The current landscape of chronic spontaneous urticaria

i. Definition and burden

Chronic spontaneous urticaria (CSU) is a condition which persists for more than 6 weeks in duration and occurs in the absence of an identifiable trigger. It tends to be more common in adults than in children and in woman than in men with peak occurrence in the third to fifth decades of life. The prevalence and incidence of CSU are not truly known. The prevalence in the general population is estimated at 0.5 to 5%, while the incidence is reported around 1.4% annually (1,2). More precise data are awaited from the Chronic Urticaria Registry (CURE) designed to improve the scientific understanding, clinical treatment and healthcare planning of chronic urticaria patients (3). An international observational study assessed a cohort of 673 adult patients with CSU whose symptoms persisted for ≥ 12 months despite treatment. Almost 50% of patients had moderate-to-severe disease activity as reported by Urticaria Activity Score (UAS) and had significant impairment in their quality of life (QoL), including significant interference with sleep and daily activities. More than 20% of patients reported ≥ 1 hour per week of missed work; productivity impairment was 27%. These effects increased with increasing disease activity (4). More than 25% of cases are resistant to H1 -antihistamines, even at higher doses, and third- and fourth-line therapies (omalizumab and ciclosporin) control the disease only in two-thirds of H1 -antihistamine-resistant patients (5). Significant healthcare resources and costs are needed to manage CSU (4,5).

ii. Phenotypes and endotypes – practical implications for management

CSU results from pathogenic activation of mast cells (MCs) and basophils, which gives rise to the release of proinflammatory mediators that support the generation of urticaria.

Activation of the high-affinity IgE receptor, Fc ϵ R1, is an important step in the development of CSU (6). This receptor is composed of an α -, β -, and two γ subunits (7). Whereas the α -subunit binds to the C ϵ 3 constant region of the IgE molecule, the β -, and γ - subunits contain cell immunoreceptor tyrosine-based activation motifs (ITAMs) which, when phosphorylated, promote activation of spleen tyrosine kinase (SYK) and downstream recruitment of a host of secondary molecules including those involved in the phosphoinositide-3 kinase (PI3K) pathway. This series of events is responsible for degranulation of mast cells and can

predispose to pathologic mast cell activation when inappropriately upregulated. SYK is recruited to the FcεR1 upon antigen stimulation, and inhibition of this protein has been shown to inhibit mast cell degranulation and production of both lipid mediators and cytokine activity (8). When mast cells from CSU patients with active urticarial disease at the time of blood sample collection were compared to those from healthy human donors, they were unsurprisingly found to release significantly more histamine in vitro than their healthy counterparts (9). Yet when these CSU patients were further subdivided into responders vs. non-responders based on their ability to degranulate in response to anti-IgE (with responders showing >10% degranulation activity), SYK levels were shown to be higher in the responder group than in the non-responder group, suggesting that this protein is a major determinant of predilection toward spontaneous degranulation. SYK expression is highly variable among the general population and is thought to correlate with the degree of IgE-mediated degranulation. Intriguingly, the presence of autoantibodies to FcεR1α or IgE do not predispose to upregulation of basophil SYK expression (10).

Accordingly, FcεRI has emerged as a viable target for the development of biologicals that act to inhibit or attenuate the activation of mast cells and basophils. At the forefront of these strategies are omalizumab, an anti-IgE monoclonal antibody reducing FcεRI surface expression, Designed Ankyrin Repeat Proteins (DARPs) inhibiting FcεRI-IgE activation through protein-protein interactions, and fusion proteins to co-aggregate FcεRI with the inhibitory FcγRIIb (11,12,13,14,15,16).

Recent evidence points towards an autoimmune aetiology in up to 50% of patients, due to MCs activating autoantibodies. Two different autoimmune mechanisms are essentially responsible for MC activation in CSU: in the majority of patients, type I autoimmunity ("autoallergy") is present, i.e., an auto-IgE-mediated immediate reaction against an autoantigen, an endogenous allergen. In a smaller proportion of CSU patients, type II autoimmunity is present, in which IgG and IgM antibodies are directed against cellular structures on MCs, for example the IgE receptor FcεRI, leading to MC activation (17,18). IgG-anti-thyroperoxidase may also be present (19,20). The presence of these autoantibodies is detected by the autologous serum skin test (ASST) and basophil tests (BTs), either basophil activation test or basophil histamine release assay (BHRA). Many CSU patients are positive in only one of these tests and they showed divergent results in about 1/3 of the cases. The authors advance the hypothesis of autologous skin signals modulating MC degranulation (21). Characterisation of this endotype in the clinic is important as it seems to be more severe and more responsive to the targeted intervention with omalizumab. The assessment of basophil FcεRI levels might also be relevant in predicting response to anti-IgE treatment (6). Patients who have a positive BHRA (~ 20% of all CSU patients) are not only more likely to have an overall more severe and prolonged CSU, but also respond significantly worse to therapy with omalizumab, an anti-IgE monoclonal antibody, which is otherwise very successful in CSU (22). Very recently, interleukin 24 (IL-24) has been identified as an IgE autoantigen in CSU (23). These patients seem to respond to autologous serum therapy (24). The epithelial derived cytokines IL-25, IL-33 and thymic stromal lymphopoietin (TSLP) have previously been shown to activate MCs and are thought to contribute to the pathogenesis of CSU (25). Both IL-4 and IL-5 contribute to the survival of MCs and they can enhance FcεRI-mediated degranulation (26, 27).

Another CSU endotype might be driven by dysregulation of MCs surface receptors or the intracellular signaling pathways within mast cells and basophils that lead to defects in trafficking or function of these cells. MCs express numerous G protein-coupled receptors (GPCRs) that lead to activation and degranulation of MCs. These include the complement C5a receptor (C5aR, CD88), to which the anaphylatoxin C5a binds, and the Mas-related G protein-coupled receptor X2 (MRGPRX2), both preferentially expressed by the skin MCs (27,28,29, 30). MRGPRX2 is a receptor that is activated by substance P, major basic protein, and eosinophil peroxidase and by various external triggers (31). A small group of MC receptors

mediates inhibitory signals. Two of these inhibitory MC receptors are Siglec-8 and CD200R (32,33).

Endothelial cells (ECs) contribute importantly to key features of CSU. Several markers of EC activation have been reported, from adhesion molecules, tissue factor, and P-selectin to D-dimers, MMP-9, endostatin, heat shock proteins, cleaved high molecular weight kininogen, and adipokines (34).

iii. *Current management*

The therapeutic goal in the treatment of CSU is to achieve complete freedom of symptoms. The current guideline for the treatment of CSU recommends second-generation non-sedating antihistamines in standard doses as the first step. If control is not adequate (after 2 – 4 weeks or earlier, if symptoms are intolerable), the antihistamines dose should be increased up to 4-fold. Individual tolerance as well as a possible increase in sedative effects at higher doses should be considered. If this does not result in sufficient control, omalizumab is additionally administered. If there is no therapeutic success after six months of treatment with omalizumab, the guidelines recommend off-label use with ciclosporin A in addition to existing therapy with H1 antihistamines. Recommended dosages are 4 mg/kg or less. In case of acute exacerbations, treatment with sufficient doses of oral glucocorticoids can be given for a short period (up to a maximum of ten days) to reduce the duration and activity of the disease. A medium-high dose of prednisolone of 20–50 mg/day for a maximum of ten days is recommended. Long-term treatment with systemic glucocorticoids or frequent “acute interventions” should be avoided at all costs due to the high rate of side effects (1,17). Management of triggering factors and of co-morbidities improves disease control. The assessment of the disease burden, defined as disease activity plus quality of life and disease control, supports the decision whether therapy is successful or should be escalated if necessary.

b. *Biologicals*

Biologic products (biologicals) include a wide range of products such as vaccines, blood and blood components, allergen vaccines, somatic cells, gene therapy, tissues, and recombinant therapeutic proteins. They are isolated from a variety of natural sources - human, animal, or microorganism - and may be produced by biotechnology methods and other cutting-edge technologies. For the purpose of this guideline we refer to monoclonal antibodies (mAb) as biologicals. In contrast to chemical compounds and small-molecule agonists or antagonists, biologicals bind a specific determinant, for example, a cytokine or receptor. Owing to this selectivity, biologicals are ideal for ‘personalised’ or ‘precision’ medicine (35).

Omalizumab inhibits the interaction between IgE and its high affinity receptor FcεRI, thus preventing mast cell and basophil activation, and blocks IgE binding to CD23 on B cells and antigen-presenting cells. The crystal structure of the complex between an omalizumab-derived Fab and IgE-Fc, with one Fab bound to each Cε3 domain was recently described. Free IgE-Fc adopts an acutely bent structure, but in the complex it is only partially bent, with large-scale conformational changes in the Cε3 domains that inhibit the interaction with FcεRI. CD23 binding is inhibited sterically due to overlapping binding sites on each Cε3 domain. Studies of omalizumab Fab binding in solution demonstrate the allosteric basis for FcεRI inhibition and, together with the structure, reveal how omalizumab may accelerate dissociation of receptor-bound IgE from FcεRI, exploiting the intrinsic flexibility and allosteric potential of IgE (36). Its use in CSU is supported by the key role of IgE and its high-affinity receptor, FcεRI, in the degranulation of skin mast cells that drives the development of the signs and symptoms of CSU, itchy wheals, and angioedema.

c. *Purpose of the EAACI Guidelines*

Delivering high-quality clinical care is a central priority for allergists, dermatologists, paediatricians, internal medicine and other specialities caring for patients with CSU. The European Academy of Allergy and Clinical Immunology (EAACI) develops and updates each year resources to help healthcare professionals (HCP) and researchers to design the best interventions, deliver high standard care and to assess their actions and decisions for purposes of quality improvement and/or reporting.

EAACI guidelines include recommendations for the management of patients with particular conditions or diseases. Guidelines are developed using a systematic process, and are based on available evidence and the clinical experience and expertise of all interested stakeholders. Following the rapid accrual of evidence for omalizumab in CSU together with an advancement of guideline development methodologies a guideline focused on the use of omalizumab in CSU was therefore needed.

The current EAACI guideline for the use of omalizumab in CSU is focussed only on treatment with omalizumab for CSU. It does not address any topics related to CSU diagnosis, concurrent treatment, or monitoring adherence.

The EAACI Guideline for the use of omalizumab in CSU is not intended to impose a standard of care. Instead, it provides the framework for rational decisions for the use of omalizumab in CSU by HCPs, patients, third-party payers, institutional review committees and other stakeholders. Statements regarding the underlying values and preferences as well as qualifying remarks accompanying each recommendation are an integral part of the Guidelines and aim to facilitate more accurate interpretation. They should never be omitted or ignored when quoting Guidelines recommendations.

i. Target audience

The target audience includes all HCPs involved in the management of CSU, patients and caregivers, basic scientists involved in biologicals development, regulatory authorities and policy makers.

ii. Biologicals included – rationale for choosing

This EAACI guideline provides recommendations for the use of omalizumab in patients with CSU. Omalizumab is currently the only biological with regulatory approval for the treatment of CSU.

Additional comments are provided for the biologicals and other targeted interventions currently tested and not yet approved and for doses/routes not approved by regulatory authorities.

II. Methods

This EAACI guideline followed the GRADE methodology (available at www.gradeworkinggroup.org). Training was conducted with all members of the guidelines development group (GDG) to prepare them for their roles, including specific sessions on the GRADE methodology.

a. The Guideline Development Group

A Core Leadership Team (table S1) supervised the project and was responsible for defining the project scope, drafting the clinical question to be addressed by the guideline, coordinating the search, and drafting the manuscript together with the Voting Panel (table S1). The project was led by three chairs with both content and methodologic expertise. The Core Leadership Team received support from a methodologist team, who advised on the process and provided input on the GRADE summary of findings (SOF) tables. The methodologist team conducted the systematic literature review (SR) for the clinical question, graded the quality of evidence, developed the SOF tables, and provided the evidence reports. Narrative reviews were conducted by different content specialist subgroups for each topic to be covered to complement the SR.

The Voting Panel, composed of content experts, decided which clinical questions are to be asked and which outcomes are critical, important and of low importance, and voted for the final recommendations after reviewing the evidence provided by the methodology team and the narrative reviews. The Voting Panel included specialists with expertise and clinical experience in treating CSU, biologists and clinical immunology experts, as well as patient representatives.

In accordance with EAACI policy, everyone who was intellectually involved in the project (i.e., considered for guideline authorship) disclosed all potential conflict of interest (COIs) in writing at the beginning, middle, and end of the project. The Guideline Oversight Committee (table S1) was responsible for developing and implementing rules related to COIs.

b. Definitions

The GDG framed the clinical question as “*Is the treatment with omalizumab efficacious and safe for patients with CSU?*” (Table 1). For the purpose of this SR, the population was defined as patients 12 years or older with a diagnosis of CSU inadequately controlled by H1-antihistamine treatment.

For the recommendations the population was defined as in the clinical trials that informed the regulatory approval.

c. Task Force question and prioritization of outcomes

Clinically relevant interventions and comparators were developed balancing comprehensiveness with feasibility (table 1). The most challenging decision in framing the question was how broadly the patients and intervention should be defined. The underlying biology of CSU suggested that across the range of patients and interventions it is plausible that the magnitude of effect on the key outcomes is different, thus the GDG defined subpopulations based on age (12-17 years old, > 18 years old) and on omalizumab dose (150 mg vs 300 mg).

As required by the GRADE approach CSU-related outcomes were prioritised in a first step by the GDG using a 1 to 9 scale (7 to 9 critical; 4 to 6 important; 1 to 3 of limited importance). The critical outcomes were weekly urticaria activity score (UAS)-7, the weekly itch severity score (ISS)-7, and safety (drug-related adverse events and serious adverse events). Important outcomes were: QoL (assessed with Dermatology Life Quality Index (DLQI) and the Chronic Urticaria Quality of Life Questionnaire (CU-Q2oL)); resource utilisation (assessed with work productivity and activity impairment (WPAI), with two sub-scales separately, work impairment score (WIS) and activity impairment score (AIS)); and rescue medication use (assessed with number of tablets of diphenhydramine per week) (Table 2). After reviewing the evidence, the prioritisation of the outcomes was reassessed to ensure that important outcomes that were not initially considered are included and to reconsider the relative importance of outcomes in light of the available evidence. All CSU-related relevant outcomes were addresses simultaneously.

The GDG also defined and addressed clinical questions not covered by the systematic review (table 3).

d. Minimal important difference

To evaluate the imprecision for each outcome their minimal important difference (MID) thresholds were considered: 9.5 to 10.0 points for UAS-7 (37,38,39,40,41,42), 4.5 to 5 points for ISS-7 (39,40,41) and 2.24 to 3.10 for DLQI (43).

e. The GRADE approach (search, appraisal of the evidence)

Key principles and provisions, key terms, descriptions, drug categories, PICO (population, intervention, comparator, and outcomes) questions, search methodology and evidence reporting used in the guideline development process were predefined.

A systematic review was conducted to inform the recommendations (44). A GRADE SOF table was provided for the PICO question. The quality of evidence was evaluated based on GRADE

quality assessment criteria by two independent reviewers and discordance resolved by consensus. Quality assessment includes the risk of bias (ROB) of included trials, the likelihood of publication bias, inconsistency between trial results, indirectness of the evidence (e.g., differences between populations, interventions, or outcomes of interest in the group to whom the recommendation applies versus those who were included in the studies referenced), and imprecision (wide confidence intervals, usually due to a small number of patients or events, or those situations where clinical decision-making would differ at the extremes of the confidence interval) (45,46,47). The quality of evidence for each outcome was rated as high, moderate, low, or very low. In the absence of any data, the level of evidence was rated as very low, based on clinical experience only. Search results were pooled in an evidence report as SOF tables and accompanied by a qualitative summary of the evidence the PICO question. The Content Panel reviewed the drafted evidence report to address evidence gaps prior to presentation to the Voting Panel.

f. Additional evidence

In support of formulated recommendations, the GDG performed narrative reviews collecting evidence on phase IV, observational, real-world trials and registries and on clinical questions not addressed by the SR (table S2).

g. Consensus building and formulating recommendations

After reviewing the evidence report and the additional evidence, the Voting Panel discussed and consented by voting in a hybrid meeting (face-to-face and online) in January 2020 on the final recommendations of this Guideline. Due to the pandemics the publication was delayed with one year. For each outcome, the Voting Panel heard an oral summary of the evidence and voted on the wording, direction and strength of the related recommendation. A 70% consensus threshold was reached for all recommendations presented below. The recommendations follow the data included in the evidence-to-decision (EtD) tables and take into consideration the balance of desirable and undesirable consequences, the quality of evidence, patients' values and preferences, feasibility, and acceptability of various interventions, use of resources paid for by third parties, equity considerations, impacts on those who care for patients, and public health impact (45,46,47). A strong recommendation was made in favour of an intervention when the GDG was certain that the desirable consequences outweighed the undesirable consequences. A conditional recommendation was provided if there were reasons for uncertainty on the benefit-risk profile, especially for low or very low quality of evidence. The underlying values and preferences played a key role in formulating recommendations. As the key target audience of this EAACI Guideline are HCPs and the patients they treat, the perspective chosen when formulating recommendations was mainly that of the HCPs and of the patient, although the health systems perspective was also evaluated, as per WHO recommendations for guidelines development (48). Recommendations are formulated separate by outcome. The recommendations formulated in this guideline should be used following the GRADE interpretation (table 4). These recommendations should be reconsidered when new evidence becomes available and an update of this guideline is planned for 2025.

Where no evidence was available the GDG formulated expert-based recommendations.

The Guideline was available on the EAACI website for two weeks (June 2021) for public comment and it underwent external peer-reviewed. All comments received were carefully reviewed by the GDG and incorporated where applicable.

h. Final review and approval of the guideline by EAACI

In addition to journal and external peer review, the EAACI Scientific Committee and Executive Committee reviewed the manuscript. These EAACI over-sight groups did not mandate that certain recommendations be made within the guideline, but rather serve as peer reviewers.

III. Key recommendations

Accumulating experience with omalizumab treatment for CSU confirmed its effectiveness and safety, by reducing reduces the signs and symptoms and burden of CSU, improving QoL, and decreasing the use of reliever medication, both in the paediatric population 12-17 years old and in adults (49,50,51,52,53,54,55,56,57,58,59).

a. Omalizumab 150 mg

The summary of the supportive evidence is presented in table S3. Recommendations for adults and the 12-17 years old paediatric population are based on the evidence-to-decision table 5.

Recommendations are formulated together for the adult and 12-17 years old population included in the SR (Box 1)

Box 1: Recommendation for omalizumab as add-on treatment in adults and in the paediatric population 12-17 years old with uncontrolled CSU

1. Omalizumab 150 mg is recommended in adults and adolescents with chronic spontaneous urticaria* uncontrolled under antihistamines to:	Reduce disease activity as reflected by UAS-7 and ISS-7	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Reduce rescue** medication	Conditional recommendation
2. Omalizumab has demonstrated a good safety profile however drug-related AEs should be periodically monitored		Conditional recommendation

* population: CSU refractory to antihistamines

** Rescue refers to „on demand”

Justification

Omalizumab 150 mg every 4 weeks did not result in a clinically meaningful reduction of disease activity as assessed by use of the UAS7 (MD -5; 95% CI -7.75 to -2.25; high certainty of evidence). It also failed to achieve a clinically meaningful reductions of ISS7 values (MD -2.15; 95%CI -3.20 to -1.10, high certainty of evidence), thus the GDG formulated a conditional recommendation for decreasing disease activity. Compared to standard of care, omalizumab 150 mg did not meaningfully reduce DLQI values (MD -1.95; 95%CI -3.06 to -0.83; moderate certainty of evidence), thus the GDG formulated a conditional recommendation for improving QoL. As it decreased the use of rescue medication with moderate certainty of evidence: MD -1.68 (95%CI -2.95 to -0.40), the GDG formulated a conditional recommendation.

As omalizumab 150 may increase the risk of drug-related AE with low certainty of the evidence: RR 1.40 (95%CI 0.63 to 3.13) the GDG formulated a conditional recommendation for a good safety profile advising periodical monitoring of AEs.

b. Omalizumab 300 mg

The summary of the supportive evidence is presented in table S4. Recommendations for adults and the 12-17 years old paediatric population are based on the evidence-to-decision tables 6 and 7.

Recommendations are formulated together for the adult and 12-17 years old population included in the SR (Box 2)

Box 2: Recommendation for omalizumab 300 mg as add-on treatment in adults and in the paediatric population 12-17 years old with uncontrolled CSU

1. Omalizumab 300 mg is recommended in adults and adolescents with chronic spontaneous urticaria* uncontrolled under antihistamines to:	Reduce disease activity as reflected by UAS-7 and ISS-7	Strong recommendation
	Improve quality of life	Strong recommendation
	Reduce rescue** medication	Conditional recommendation
	Decrease resource utilisation	Conditional recommendation
2. Omalizumab has demonstrated a good safety profile however drug-related AEs and SAEs should be periodically monitored		Conditional recommendation

* population: CSU refractory to antihistamines

** Rescue refers to „on demand

Justification

Omalizumab 300 mg every 4 weeks led to a clinically meaningful decrease in UAS-7 (MD -11.05; 95% CI -12.87 to -9.24) and in the ISS-7 (MD -4.65; 95%CI -5.41 to -3.89), both with moderate certainty of evidence, thus the GDG formulated a strong recommendation for decreasing disease activity. It also DLQI values (MD -4.01; 95%CI -4.94 to -3.08), with an improvement in QoL above the MID with high certainty of evidence. This was paralleled by a significant decrease in CU-Q2oL (MD -15.34; 95%CI -24.84 to -5.84), consequently a strong recommendation for improving QoL was formulated. There was moderate certainty for reducing rescue medication (MD -2.04 (95%CI -3.19 to -0.88)), thus a conditional recommendation was formulated. Compared to standard of care, omalizumab 300 mg improved WIS (MD -24.24; 95%CI -35.74 to -12.74) and AIS (-26.59; 95% CI -37.36 to -15.72) at 24 weeks (58). However, as data come from only one RCT a conditional recommendation was formulated

Omalizumab 300 mg slightly increased drug-related AEs (RR 1.37 (95%CI 0.67 to 2.82) with low certainty and decreased with moderate certainty drug-related SAEs (RR 0.77; 95% CI 0.20 to 2.91)), although results are inconclusive due to the small RR. One study reported a single anaphylactic episode during the open-label phase of the study (57). The GDG formulated a conditional recommendation for a good safety profile advising periodical monitoring of AEs and SAEs.

c. Subgroups: stratified by co-morbidities

The GDG evaluated the evidence for omalizumab efficacy in CSU associated with other co-morbidities not included in the SR (table S2) and formulated a conditional recommendation, expert opinion based on the efficacy of omalizumab in patients with CSU and other co-morbidities (box 3).

Box 3: recommendation for omalizumab in adults and 12-17 years old patients with both CSU associated with other co-morbidities

Omalizumab may be of particular benefit in adults and 12-17 years old patients with both CSU associated with other co-morbidities (chronic inducible urticaria, allergic asthma, allergic rhinitis)	Conditional recommendation, expert opinion based
---	--

d. Biomarkers predicting response

Free (non-Omalizumab bound) IgE levels in serum are a measure of effective omalizumab dosing. Recent findings point to a possible role of total IgE as a marker of CSU disease activity, endotypes, and responses to treatment (60,61,62,63). A recent review of 141 publications showed that up to 50% CSU patients had elevated total IgE serum levels, but normal or very low total IgE levels also occurred. High total IgE may represent high disease activity, longer disease duration, high chance of responding to omalizumab treatment, quick relapse after stopping omalizumab, and lower chance of responding to cyclosporine. Low IgE, in contrast, may suggest Type IIb autoimmune CSU, poor response to treatment with omalizumab and a better chance to benefits from cyclosporine treatment. Furthermore, IgE in different CSU cohorts may have different physicochemical properties that could explain differences in treatment responses to IgE-directed therapies (64).

In support of the EAACI guideline recommendation a SR was performed, including all published studies evaluating the following predictive biomarkers for omalizumab efficacy: IgE and IgG autoantibodies to high- and low-affinity IgE receptors (FcεRI and FcεRII), total IgE levels, thyroid autoimmunity (IgE and IgG autoantibodies anti antithyroperoxidase and/or anti thyroglobulin), eosinopenia, basopenia, eosinopenia associated with basopenia, IgE and IgG autoantibodies to tissue factor, autologous serum skin test, basophil activation test, basophil histamine release assay positivity, high D-dimer, high CRP, high ESR, antinuclear antibodies (tables S5-S42 and figures S1-S6). For all biomarkers evaluated the evidence is very uncertain. However, the CDG formulated a conditional recommendation, with low level of evidence, for high total IgE and stated that the other biomarkers evaluated need further exploration in prospective trials with prediction of response as primary end-point.

Box 4: recommendation for biomarkers in prescribing omalizumab in adults and 12-17 years old patients with CSU

High total IgE may indicate a higher chance of responding to omalizumab treatment and of quick relapse after stopping omalizumab.	Conditional recommendation, low level of evidence
---	---

e. Implementation considerations

For omalizumab 300 mg every 4 weeks the GDG formulated strong recommendations for the reduction in disease activity and for improving QoL and conditional recommendations for reducing rescue medication and resource use and for safety-related outcomes. For omalizumab 150 mg every 4 weeks only conditional recommendations were formulated for all outcomes. According to GRADE for strong recommendations most individuals should receive the intervention and the recommendation can be adapted as policy or performance measure in most situations (table 4).

However, the GDG cautions on several unsolved key pillars supporting the implementation of these recommendations, such as independent high-quality cost-effectiveness studies,

selection of responders, documentation of the disease modifying effect together with long-term safety data, studies addressing a priori CSU together with its co-morbidities. The cost-effectiveness of omalizumab based on real-world treatment patterns is unknown. Including broader evidence on treatment discontinuation, caregiver burden, and rescue medication and resource use reduction from real-world studies and CSU registries may better reflect the effects and value of biologicals for all healthcare stakeholders (3). Last but not least the value of the recommendations depends also on the setting in which the current guideline will be implemented, as recommendation suitable for resource-rich environments might change from strong to conditional in resource-poor environments (Box 5).

Box 5: Factors impacting the implementation of recommendations for the use of omalizumab in CSU (adults and 12-17 years old)

1. Cost-effectiveness, especially independent real-world evidence
2. Long-term safety data
3. Immune modulation/disease modifying effect
4. Stratification* based on biomarkers**
5. Patient's preference
6. Availability of resources

* Stratification – safety and efficacy

**Biomarkers include both clinical and laboratory features

IV. Practical approach

a. Definition of response; continuation and stopping rules

Disease control is a major treatment aim in CSU. UAS-7 has become the standard for assessing CSU disease activity in clinical studies and the routine management of patients, but it does not evaluate disease control. The urticaria control test (UCT) is a 4-item questionnaire about physical symptoms, QoL, treatment effects, and urticaria control over the previous 4 weeks (65). Responder rates using UCT score of greater than or equal to 12 and UAS value of 6 or less yielded very similar results (66). Unfortunately, there are no validated cut-offs within these tools used to define response, nor an agreement on the interval when activity and control should be evaluated after the initiation of omalizumab treatment. Approximately 30% of patients remain symptomatic at licensed doses of omalizumab 150 mg and 300 mg, even after a treatment period of over 6 months. In the recent years, there have been several studies on up-dosing omalizumab_[AG1], suggesting that the individualized approach for urticaria treatment with omalizumab is useful. Real-world data mainly show that up-dosing/dose adjustment evaluated with the assessment of disease activity (UAS-7) and control (UCT) achieves better clinical response to omalizumab with a good safety profile in a subgroup of patients with CSU (67).

Box 6: Recommendations for practical use of omalizumab in adults and 12-17 years old patients with CSU

The evaluation of response should be done after 4-6 months of treatment	Conditional recommendation, expert opinion based
As there are no validated criteria for defining response to omalizumab in CSU the GDG recommends a composite end-point combining evaluation of disease activity (UAS-7 and ISS-7), disease control (UCT), and with other measures of QoL	Conditional recommendation, expert opinion based

A pre-established cut-off reached through shared decision making with the patient should be used	Conditional recommendation, expert opinion based
Stopping omalizumab should be considered if a significant AE occurs	Conditional recommendation, expert opinion based
For insufficient response up-dosing may be considered	Conditional recommendation, low quality evidence

In addition, adherence to background treatment and to avoidance of CSU triggers measures should be evaluated before deciding to stop omalizumab due to lack of efficacy.

b. Monitoring treatment
i. Adverse events

Use of omalizumab is associated with reported side effects ranging from local skin inflammation at the injection site to systemic anaphylaxis. Omalizumab binds to the constant region of free IgE only and, therefore, does not cause mast cell degranulation. However, omalizumab has been reported to cause anaphylaxis in <0.1% of patients, with reactions being delayed in some cases (68,69,70). To date, the mechanisms through which omalizumab induces adverse reactions are still unknown. Treatment with omalizumab results in a markedly increased sensitivity of basophils to IgE-mediated stimulation in terms of the number of IgE molecules required to produce a given response (71). In a recent experimental model, it was shown that omalizumab can induce skin inflammation and anaphylaxis by engaging FcγRs, and demonstrated that Fc-engineered versions of the mAb could be used to reduce such adverse reactions without compromising efficacy (72). A non-IgE-mediated anaphylactic response to the polysorbate excipient might also be involved (73).

In pre-marketing clinical trials in patients with asthma, anaphylaxis was reported in 3 of 3,507 (0.1%) patients. In post-marketing spontaneous reports, the frequency of anaphylaxis attributed to omalizumab use was estimated to be at least 0.2% of patients based on an estimated exposure of about 57,300 patients from June 2003 through December 2006. The risk factors for these anaphylactic reactions were uncertain given the limitations of spontaneous reports and the lack of control data. Due to this increase in the rate of anaphylaxis, a black box warning was added to the omalizumab label in 2007. To understand the risk factors associated with anaphylaxis among omalizumab-treated patients, a pharmacosurveillance data repository (Q4458g, X-PAND) was initiated in early 2009 as a post-marketing commitment. Data collected included clinical histories, immunogenicity assessment, and an optional allergy skin test. Thirty cases of anaphylaxis and 120 controls were considered to give 79% power to detect a 4-fold increase in the risk for an anaphylactic reaction assuming a 10% prevalence for an identified risk factor and a 5% type I error (2-tailed). Prespecified potential clinically meaningful risk factors (eg, presence of food allergy, preomalizumab IgE levels, asthma severity, sex, and age) were assessed for effect on the frequency of anaphylaxis. Most cases of anaphylaxis (24 of 30; 80%) included symptoms categorized as cutaneous/subcutaneous/mucosal (ie, lips, tongue, palate, uvula) and respiratory (ie, nose, laryngeal, lung). Most (70.0%) events occurred within 1 hour of omalizumab dosing (only 1 event occurred after 2 hours [ie, at 3.5 hours]). Median (range) time from the last dose of omalizumab to the anaphylactic event was 30 (0-210) minutes. Eleven of 28 (39.3%) patients in whom the number of previous doses had been recorded

experienced anaphylaxis within the first 3 doses of omalizumab; anaphylaxis occurred after 4 to 20 doses in 8 (28.6%) cases and after more than 20 doses in 9 cases (1 case after >60 injections). None of the anaphylactic events resulted in disability or death. Anaphylactic events were considered life-threatening in 12 of 30 (40.0%) cases and required hospitalization in another 6 (20.0%) cases. Treatment of anaphylaxis included the use of antihistamines (23 of 30; 76.7%), epinephrine (21 of 30; 70.0%), systemic corticosteroids (19 of 30; 63.3%), and inhaled β -agonists (13 of 30; 43.3%). Bivariate conditional logistic regression analysis indicated that among omalizumab users, a history of anaphylaxis to food, medication, or other causes increased the subsequent risk of anaphylaxis associated with omalizumab use (OR, 8.1; 95% CI, 2.7-24.3). The US Food and Drug Administration examined this information and advised including it in an updated Xolair US package insert. Total number of omalizumab doses, food allergies, female sex, presence of urticaria/hives, and race also were identified as potential risk factors for anaphylaxis associated with omalizumab. The absolute risks for anaphylaxis may be estimated on the basis of overall risks of 0.1% (based on clinical trial data) or 0.2% (based on post-marketing reports) of users. Assuming an overall 0.2% risk, the absolute increase in risk can be indirectly estimated from the OR, and it would be approximately 0.62% for patients with a history of anaphylaxis and 0.08% for patients with no history of anaphylaxis, resulting in a risk difference (ie, attributable risk) of 0.54% (74). As part of this study, an assay that could detect antibodies of IgE isotype to omalizumab was developed. Using this assay there was no apparent correlation between either anaphylaxis or skin test reactivity and the presence of antibodies of IgE isotype to omalizumab (75). As a result of this study FDA withdraw the black-box warning. Furthermore, as the reported incidence of anaphylaxis continues to be low, in several countries omalizumab became licensed for home administration.

Box 7: Recommendations for managing anaphylaxis under omalizumab treatment for CSU

The occurrence of anaphylaxis following treatment with omalizumab in CSU is an event of special interest that should be reported appropriately in order to improve the post-marketing surveillance data	Conditional recommendation, expert opinion based
The first 3 doses should be administered in a setting with experience in managing anaphylaxis; an observation period of 30 minutes post-administration is recommended	Conditional recommendation, expert opinion based
Consultation with an allergist is encouraged if risk factors for anaphylaxis are present	Conditional recommendation, expert opinion based
As most cases are mild/moderate and respond well to anaphylaxis treatment omalizumab should not be discontinued	Conditional recommendation, expert opinion based
Home administration is an option starting with the 3 rd dose with the condition that the patient has been provided with an anaphylaxis action plan and proper education	Conditional recommendation, expert opinion based

While elevated serum IgE is generally associated with allergic/atopic conditions, very low or absent IgE may hamper anti-tumour surveillance, indicating the importance of a balanced IgE-mediated immune function (76). The Epidemiologic Study of Xolair Evaluating Clinical Effectiveness and Long-Term Safety in Patients with Moderate to Severe Asthma (EXCELS), a 5-year observational cohort study conducted in patients 12 years or older with moderate-to-severe asthma to evaluate the long-term safety of omalizumab, primarily the risk of

malignancy (77). The authors conclude that the results “suggest that omalizumab is not associated with an increased risk of malignancy.” However, the potential for unmeasured/uncontrolled confounding, the selection biases introduced by the enrolment of “prevalent users” (previously exposed to omalizumab) and the initial exclusion of patients with a history of cancer or a premalignant condition, and the high study discontinuation rate significantly limit the ability of the study to rule out a malignancy risk with omalizumab treatment. A systematic review and meta-analysis of intervention and observational studies evaluated whether prolonged treatment with omalizumab influences development or progression of solid epithelial cancer in patients with atopic asthma or CSU. Only 12 studies reported outcomes of interest and none included CSU. There was insufficient evidence to determine whether long-term treatment with omalizumab influences development or progression of solid epithelial cancer in these patient populations (78). Of note, EXCELS showed a higher rate of cardiovascular and cerebrovascular events (79). [AG2]

ii. Routine laboratory monitoring

Box 8: Recommendation for routine laboratory monitoring for omalizumab treatment for CSU

No routine laboratory monitoring is recommended for omalizumab in CSU	Conditional recommendation, moderate quality evidence
---	---

iii. Infections and response to vaccination

Omalizumab reduces inflammation by blocking proinflammatory cytokines and may even have antiviral effects. It affects mast cells, blocking the release of inflammatory agents such as histamine and protease in addition to proinflammatory cytokines including IL-1, IL-6, and IL-33 and has been shown to increase antiviral immunity through downregulation of the high-affinity IgE receptor on plasmacytoid dendritic cells, which is essential for antiviral immune responses (80). In inner-city asthmatic children aged 6 to 17 years preseasonal omalizumab treatment prevented the viral exacerbations in the fall, while increasing IFN- α responses to rhinovirus (81). Treatment with omalizumab was effective and safe in patients with ABPA, regardless of comorbid chronic respiratory tract infections (chronic *Pseudomonas aeruginosa* or nontuberculous mycobacterial infection of the lower respiratory tract) (82). There was no safety signal from RCTs or RWE data on the use of omalizumab and increased risk of bacterial, viral or fungal infections, nor on the use of anti-infectious vaccines.

Although the role of IgE in immunity against helminth parasites is unclear, there was concern that omalizumab may be unsafe in subjects at risk of helminth infection. In an exploratory study of allergic subjects at high risk of helminth infections, omalizumab therapy appeared to be safe and well tolerated, but may be associated with a modest increase in the incidence of geohelminth infection (OR adjusted for study visit, baseline infection status, gender and age = 2.2 (0.94-5.15); one-sided P=0.035). Infection severity and response to helminthics appeared to be unaffected by omalizumab therapy (83). The European Society of Clinical Microbiology and Infectious Diseases recommends that pre-treatment screening for *Strongyloides stercoralis* and other geohelminths should be considered in patients who come from areas where these are endemic who are receiving IgE-targeted agents (84).

Box 9: Recommendation for management of infections and vaccinations under omalizumab treatment for CSU

Omalizumab should not be discontinued in case of cutaneous or non-cutaneous bacterial, viral or fungal infections or in case a vaccination is required, however, an unexpected outcome such as serious infection or	Conditional recommendation, moderate quality evidence
---	---

vaccination failure should be reported appropriately in order to improve the post-marketing surveillance data	
Pre-treatment screening for geohelminths is recommended for patients where this infection is endemic	Conditional recommendation, expert opinion based
A 7-day window between the vaccine and omalizumab administration is recommended to unequivocally assign adverse events to either of the interventions is recommended	Conditional recommendation, expert opinion based

V. Other biologicals and small molecules tested for CSU

a. Targeting the IgE pathway: ligelizumab, quilizumab, GI-310

A high-affinity monoclonal anti-IgE antibody, ligelizumab, has recently been developed to overcome some of the limitations associated with the clinical use of omalizumab. Ligelizumab shows superior inhibition of IgE binding to FcεRI, basophil activation, IgE production by B cells and passive systemic anaphylaxis in an in vivo mouse model. However, ligelizumab was less potent in inhibiting IgE:CD23 interactions than omalizumab (85). Overall, ligelizumab, has ~ 50 times higher affinity for IgE than omalizumab. The results of a recently published multicenter, randomized, controlled phase II study show that ligelizumab is a highly effective therapy for CSU, with a higher rate of complete responders as compared with omalizumab, and with a very rapid and effective response and a longer lasting effect (86). In the SR conducted for the EAACI guidelines the certainty of evidence for ligelizumab in decreasing UAS-7 was categorised as low as the decrease was below the MID (-2,28; 95%CI -7,72 to -3,16). However, ligelizumab showed a good safety profile as it may decrease drug-related adverse events (risk ratio 0.72; 95%CI 0.50 to 1.05) (table S43). Phase 3 trials in adults and adolescents with CSU are currently ongoing and have yet to confirm these results (NCT03580369, NCT03580356).

Quilizumab, a [AG3]humanized, afucosylated, monoclonal IgG1 antibody, binds membrane IgE at the M1-prime segment, which is absent in soluble IgE. In animal studies, quilizumab bound membrane IgE on IgE-switched B cells and plasmablasts and depleted them through apoptosis and antibody-dependent cell-mediated cytotoxicity (87). In the QUAIL study (NCT01987947), although quilizumab reduced median serum IgE level approximately 30% in patients with CSU, it did not cause clinically meaningful improvements in the ISS-7 or the UAS-7 (88). A similar failure to achieve a significant clinical effect was encountered in asthma (89) and the drug was discontinued.

GI-301 (GI-Innovation) is a novel long-acting IgE trap-Fc fusion protein that, like omalizumab and ligelizumab, binds circulating IgE. Similar to ligelizumab GI-3012 has higher and more durable binding to IgE than omalizumab (90).

b. Targeting the T2 pathway: dupilumab, benralizumab, mepolizumab, reslizumab

Recently, a small case series has now demonstrated that dupilumab, an anti-IL-4Rα antibody, is effective in adult patients with CSU (91). The efficacy of dupilumab in urticaria is currently being investigated in several clinical trials, both in CSU and cholinergic urticaria (NCT03749135, NCT03749148, NCT04180488).

Mepolizumab and reslizumab have been successfully used in the treatment of individual patients with CSU (92,93). Positive results of smaller controlled trials with benralizumab were recently published (94,95). Benralizumab and mepolizumab are currently in clinical trials to test their efficacy in CSU (NCT04612725, NCT03494881).

c. Targeting mast cell receptors

Lirentelimab, an anti-Siglec-8 monoclonal antibody, was recently shown to inhibit MC activation and lead to extensive depletion of eosinophils. Lirentelimab has been successfully tested in an open-label phase IIa pilot study in patients with omalizumab-naïve and omalizumab-refractory CSU, as well as in patients with symptomatic dermographism or cholinergic urticaria (96). However, larger controlled studies to confirm the safety and efficacy of the drug in the treatment of CSU are still pending. Initial findings on the safety of lirentelimab were gathered in a study on eosinophilic gastritis and duodenitis (97).

The monoclonal antibody LY3454738 directed against CD200R is currently tested in a randomized, controlled phase 2 trial in patients with CSU (NCT04159701).

Preliminary results from a phase 1 study in 32 healthy volunteers with the anti-Kit antibody CDX-0159 indicate that treatment leads to a substantial reduction of MCs. A single intravenous dose-dependent administration led to an almost complete reduction of basal tryptase in the blood after only a few days. In the two higher doses, there was a sustained suppression of tryptase until the end of the observation period of 71 days (98). CDX-0159 is currently in ongoing clinical trials for CSU. [AG4]

VI. Discussion

a. Relevance of the EAACI guideline in relation to existing CSU guidelines

The EAACI Guidelines recommendations for the use of omalizumab in CSU are formulated per outcome and per dose. The GRADE approach was used to rate the certainty of the evidence. The outcomes included were prioritised beforehand and the minimal important difference was considered when available for all CSU-related outcomes. Besides judging the risk of bias, the recommendations considered all relevant aspects related with the certainty of evidence like heterogeneity, indirectness or imprecision of the results. A critical appraisal of the evidence not included the SR provided additional support for the GDG in formulating recommendations. The recommendations follow the data included in the evidence-to-decision tables and take into consideration the balance of desirable and undesirable consequences, quality of evidence, cost-effectiveness, patients' values and preferences, feasibility, and acceptability of various interventions, use of resources paid for by third parties, equity considerations, impacts on those who care for patients, and public health impact

The EAACI/GA²LEN/EDF/WAO guideline for the definition, classification, diagnosis and management of urticaria recommends adding on omalizumab 300 mg for the treatment of patients with CU unresponsive to 2nd-generation H1-antihistamines, mentioning that omalizumab has high-quality evidence, high cost, very good safety profile, and very good efficacy. It also acknowledges the preventive effect on angioedema in CSU (1). However, the recommendation is formulated on the basis of >90% consensus, and there are no separate recommendations for different outcomes (disease activity, QoL, rescue use, safety), nor a strength of the recommendation. EAACI guidelines also offer a conditional recommendation for the 150 mg dose.

The 2014 Joint Task Force on Practice Parameters mentions omalizumab (together with cyclosporine) under Annotation 4 (Add an immunosuppressant or biologic agent) as having the greatest published experience documenting efficacy in patients with CU compared with all other alternative agents. Besides being recommended in step 4, together with immunosuppressants, there is no other specific recommendation in CSU besides the fact that omalizumab should be considered for refractory CU if this is favourable from the standpoint of balancing the potential for benefit with the potential for harm/burden and cost and the decision to proceed is consistent with patients' values and preferences (99).

The Italian Society for Pediatrics, the Italian Society for Allergy and Immunology, and the Italian Society for Pediatric dermatology convened a multidisciplinary panel that prepared clinical guidelines for diagnosis and management of chronic urticaria in childhood (100). The panel recommends omalizumab in children 12 years of age and older with CSU added to second-generation H1-antistamines as a second-line therapy when second-generation H1-antistamines alone do not give adequate relief. (Level of evidence I. Strength of recommendation A). Again, there are no separate recommendations per outcome or per dose.

The KAAACI/KDA Evidence-Based Practice Guidelines for CSU in Korean Adults and Children recommends omalizumab for patients with CSU that do not respond to H1-antihistamines (strong recommendation, moderate quality of evidence) (101). The guidelines follow the GRADE approach and are based on a SR conducted specific for these guidelines but similar to the previous guidelines is does not provide specific recommendations per outcome or per dose. However, these guidelines offer an additional recommendation for omalizumab for patients with CU is not controlled with H1-antihistamines and immunomodulators (conditional recommendation, very low quality of evidence).

b. Future perspectives, barriers and facilitators

i. Precision medicine using endotyping and multiple upstream targets

The separation into type I and type IIb autoimmunity and further validation of biomarkers is of utmost importance to select responders to omalizumab (17,22,102,103,104). In addition, a better understanding of the “non-canonical” mechanisms of action of omalizumab, such as effects on mast cell releasability or the coagulation cascade or targeting membrane-IgE in IgE+ B-cells, reducing IL4R expression and IgE synthesis and decreasing the number of these cells, possibly by causing B-cell anergy, should be further prioritised (13, 105, 106). Targeting the epithelial derived cytokines or the GPCRs or MCs depletion are attractive pathways for endotype-driven management of CSU (107).

ii. The disease modifying effect

The “holy Grail” for the use of biologicals in CSU is to validate their disease modifying potential. One RCT assessed worsening in UAS7 (≥ 12 points for ≥ 2 consecutive weeks) DLQI worsening (≥ 3 points increase) for omalizumab 300 mg after stopping the treatment (58). Patients in the placebo group were at higher risk for UAS7 worsening (RR 2.88; 95%CI 1.79 to 4.63) and had higher likelihood of DLQI worsening (RR 3.34;95% CI 2.07 to 5.40) as compared to the omalizumab 300 mg group. This result might indicate an “expanded” effect of omalizumab, however currently omalizumab did not demonstrate a convincing disease modifying effect in CSU, as the efficacy is lost a few weeks or months after the treatment is stopped.

iii. Long-term safety

Omalizumab has evidence for long-term safety above 5 years for the treatment of severe asthma, both in the adult and the paediatric population (108, 109,110). However, there are no data on the long-term use in CSU. Thus, postmarketing surveillance, especially collected through structured registries like CURE is of utmost importance

iv. Considerations for the pediatric population

Data on the efficacy and safety of omalizumab in the 12-17-years old CSU patients are limited and evidence for long-term use (> 1 year) is lacking. The development of new drugs for the treatment of paediatric CSU proves difficult due to the limited availability of a very heterogenous population to enter randomised placebo-controlled trials in combination with the stringent requirements of the Paediatric Investigational Plan (EMA) or Paediatric Study Plan

(FDA). Registries and large-scale international consortia evaluating paediatric CSU could help to overcome this major unmet need in the field of omalizumab for CSU.

v. Efficacy versus effectiveness in a real-world setting

Several retrospective real-life cohorts and CSU registries report similar impact of omalizumab on CSU severity as in RCTs with an acceptable safety profile (table S2).

vi. Cost-effectiveness

To determine the cost-effectiveness of omalizumab relative to standard of care (up to four times the daily dose of H1 -antihistamines) in the Netherlands from a societal perspective a Markov model used consisted of five health states based on UAS- 7 with a 10-year time horizon. The incremental cost-effectiveness ratio (ICER) of omalizumab versus standard of care was €17 502 per quality-adjusted life-year (QALY) gained. Productivity costs played an important role in the value of the ICER as by discarding productivity costs resulted in an ICER of €85 310 per QALY (111). A similar model applied in UK showed a deterministic ICER of £3183 in the base case, implying that omalizumab was associated with increased costs and benefits relative to standard of care (112). Both studies were considered by the GDG in the EtD tables for formulating recommendations.

vii. Additional major unmet needs and research priorities

The GDG proposed several key areas of interest both for the clinician and the basic researcher and from the health-care point of view (box 10). Unmet needs have been assessed from the perspectives of different stakeholders.

Box 10: Gaps in evidence for the use of omalizumab in CSU and plan to address		
Gaps in evidence	Plan to address	Priority
Standardising the use in clinical practice <ol style="list-style-type: none"> 1. Criteria for responders and suboptimal response (early stopping rules) 2. Switching rules 3. Duration of treatment in responders (late stopping rules) 4. Long-term treatment regimen in responders: longer interval, down-dosing, possibility of stopping treatment, switch to strategies like topical application, etc. 5. Identification of factors related to failure 6. Routine measurement of ADA 	Prospective trials testing the clinical question followed by validation in independent population	High
Implementation of guidelines for the use of biologicals in clinical practice	In-depth education of HCPs on CIU pathogenic mechanisms and in recognising the endotype	High
Improving evaluation by combining clinical and molecular outcomes	Multidimensional endotyping validating skin and systemic biomarker profiles	High
Long-term safety data (> 5 years) ^[AG5]	Well-structured post-marketing surveillance using CSU registries	High
Assess the long-term efficacy/disease modifying effect (after treatment cessation)	Identify biomarkers related to the course of CSU Well-designed RCT and real-life studies focusing on long-term efficacy	High

	Mechanistic studies at a single cell level	
Efficacy and safety data in the paediatric population	RCT and RWE trials/registries focused primarily on the paediatric population	High
Efficacy and safety in selected populations (pregnancy,) and in high-risk populations	RCT and RWE trials/registries focused primarily on these populations	High
Cost-effectiveness	Sectoral and generalised cost-effectiveness analysis, including the real-world perspective Long-term perspective as disease modifying intervention and thereby influence long-term cost	High
Use of biomarkers for stratification	Proof of concept studies evaluating patient selection based on biomarkers	High
Impact of multi- morbidities	Studies evaluating the global effect of biologicals on multi-morbidities	High
Fair accessibility to CSU correct diagnosis and optimal targeted treatment	Reorganisation of CIU care Implementation of the patients' perspective from research to models of care Implementation of management pathways/clinical decision systems	High
Comparison between biologicals available for CSU (approved and currently tested)	Independent head-to-head comparison between biologicals, ideally with cross-over design	High
Alignment of studies (including RWE) with guidance from regulatory bodies.	Work in partnership with regulatory bodies to continuously review trial methodology and outcomes.	Medium
The impact of age/race/ethnicity on the short and the long-term effects (efficacy and safety)	Well-designed RCT, example for personalised medicine	Medium
Does 'resistance' occur as in antibiotic or anti-cancer therapy and what are the underlying molecular mechanisms?	Well-designed RCT, example for personalised medicine	Medium
Validation of different regimens: shorter or longer intervals ('pulse-wise') rather than as a chronic ('maintenance') therapy (e.g. to prevent [AG6]resistance)?	RCTs and real-life studies testing different approaches in terms of dose, duration and route	Medium
Combination of omalizumab with other immune modulation interventions (eg small molecules)[AG7]	RCTs and real-life studies	Medium

VII. Conclusion

The addition of omalizumab for the treatment of patients with chronic spontaneous urticaria not controlled by antihistamines is supported by improved understanding of disease mechanisms and has proved so far efficacious and safe in adults and the 12-17 years old population.

There are several critical points that need further evaluation, from the effectiveness in real-world settings to the sustainability by the healthcare systems, especially if long-term administration is warranted.

This EAACI Guideline on the use of omalizumab for chronic spontaneous urticaria offer a desk reference tool for healthcare providers, patients, regulators and healthcare systems based on a critical appraisal of the current evidence and a structured approach in formulating recommendations in alignment with the key principles of personalised medicine and implementation science.

References

1. Zuberbier T, Aberer W, Asero R, et al. The EAACI/GA²LEN/EDF/WAO guideline for the definition, classification, diagnosis and management of urticaria. *Allergy*. 2018; 73(7):1393-1414
2. Fricke J, Ávila G, Keller T, et al. Prevalence of chronic urticaria in children and adults across the globe: Systematic review with meta-analysis. *Allergy*. 2020;75(2):423-432
3. Weller K, Giménez-Arnau A, Grattan C, et al. The Chronic Urticaria Registry: rationale, methods and initial implementation. *J Eur Acad Dermatol Venereol*. 2021;35(3):721-729
4. Maurer M, Abuzakouk M, Bérard F, et al. The burden of chronic spontaneous urticaria is substantial: Real-world evidence from ASSURE-CSU. *Allergy*. 2017;72(12):2005-2016
5. Gonçalo M, Giménez-Arnau A, Al-Ahmad M, et al. The global burden of chronic urticaria for the patient and society. *Br J Dermatol*. 2021;184(2):226-236
6. Deza G, March-Rodríguez A, Sánchez S, et al. Relevance of the Basophil High-Affinity IgE Receptor in Chronic Urticaria: Clinical Experience from a Tertiary Care Institution. *J Allergy Clin Immunol Pract*. 2019;7(5):1619-1626
7. Turner H, Kinet JP. Signalling through the high-affinity IgE receptor Fc epsilonRI. *Nature*. 1999; 402(6760 Suppl.):B24–30
8. Rossi AB, Herlaar E, Braselmann S, Huynh S, Taylor V, Frances R, et al. Identification of the Syk kinase inhibitor R112 by a human mast cell screen. *J Allergy Clin Immunol*. 2006; 118:749–55.
9. Saini SS, Paterniti M, Vasagar K, et al. Cultured peripheral blood mast cells from chronic idiopathic urticaria patients spontaneously degranulate upon IgE sensitization: relationship to expression of Syk and SHIP-2. *Clin Immunol*. 2009; 132:342–8.
10. MacGlashan D. Auto-antibodies to IgE and FcεRI and the natural variability of SYK expression in basophils. *J Allergy Clin Immunol*. 2018 143:1100–7
11. Egel A, Baravalle G, Hobi G, et al. Accelerated dissociation of IgE-FcεRI complexes by disruptive inhibitors actively desensitizes allergic effector cells. *J Allergy Clin Immunol*. 2014 133:1709–19
12. Davies AM, Allan EG, Keeble AH, et al. Allosteric mechanism of action of the therapeutic anti-IgE antibody omalizumab. *J Biol Chem*. (2017) 292:9975–87
13. Kaplan AP, Giménez-Arnau AM, Saini SS. Mechanisms of action that contribute to efficacy of omalizumab in chronic spontaneous urticaria. *Allergy*. 2017;72(4):519-533
14. Baumann MJ, Egel A, Amstutz P, Stadler BM, Vogel M. DARPins against a functional IgE epitope. *Immunol Lett*. 2010; 133:78–84
15. Egel A, Buschor P, Baumann MJ, et al. Inhibition of ongoing allergic reactions using a novel anti-IgE DARPIn-Fc fusion protein. *Allergy* 2011; 66:961–8
16. Zellweger F, Gasser P, Brigger D, et al. A novel bispecific DARPIn targeting FcγRIIB and FcεRI-bound IgE inhibits allergic responses. *Allergy* 2017; 72:1174–83.
17. Maurer M, Eyerich K, Eyerich S, et al. Urticaria: Collegium Internationale Allergologicum (CIA) Update 2020. *Int Arch Allergy Immunol*. 2020; 181(5):321-333
18. Konstantinou GN, Asero R, Ferrer M, et al. EAACI taskforce position paper: evidence for autoimmune urticaria and proposal for defining diagnostic criteria. *Allergy*. 2013;68:27–36
19. Schoepke N, Asero R, Ellrich A, et al. Biomarkers and clinical characteristics of autoimmune chronic spontaneous urticaria: Results of the PURIST Study. *Allergy*. 2019;74(12):2427-2436.

20. Kolkhir P, Metz M, Altrichter S, Maurer M. Comorbidity of chronic spontaneous urticaria and autoimmune thyroid diseases: A systematic review. *Allergy*. 2017;72(10):1440-1460
21. Baumann K, Marcelino J, Skov PS, et al. Autologous serum skin test reactions in chronic spontaneous urticaria differ from heterologous cell reactions. *J Eur Acad Dermatol Venereol*. 2021 Jan 21. doi: 10.1111/jdv.17131. Epub ahead of print.
22. Gericke J, Metz M, Ohanyan T, et al. Serum autoreactivity predicts time to response to omalizumab therapy in chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2017; 139(3):1059-1061
23. Schmetzer O, Lakin E, Topal FA, et al. IL-24 is a common and specific autoantigen of IgE in patients with chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2018;142:876–882
24. Yu L, Buttgereit T, Stahl et al. Immunological effects and potential mechanisms of action of autologous serum therapy in chronic spontaneous urticaria. *J Eur Acad Dermatol Venereol*. 2019;33(9):1747-1754.
25. Kay AB, Clark P, Maurer M, Ying S. Elevations in T-helper-2-initiating cytokines (interleukin-33, interleukin-25 and thymic stromal lymphopoietin) in lesional skin from chronic spontaneous ('idiopathic') urticaria. *Br J Dermatol*. 2015; 172(5):1294-30
26. Ochi H, De Jesus NH, Hsieh FH, Austen KF, Boyce JA. IL-4 and -5 prime human mast cells for different profiles of IgE-dependent cytokine production. *Proc Natl Acad Sci U S A*. 2000 ; 97(19):10509-13.
27. Soruri A, Grigat J, Kiafard Z, Zwirner J. Mast cell activation is characterized by upregulation of a functional anaphylatoxin C5a receptor. *BMC Immunol*. 2008;9:29
28. Elieh Ali Komi D, Shafaghat F, Kovanen PT, Meri S. Mast cells and complement system: Ancient interactions between components of innate immunity. *Allergy*. 2020;75(11):2818-2828
29. Fujisawa D, Kashiwakura J, Kita H, et al. Expression of Mas-related gene X2 on mast cells is upregulated in the skin of patients with severe chronic urticaria. *J Allergy Clin Immunol*. 2014; 134(3):622-633
30. Shtessel M, Limjunyawong N, Oliver ET, et al. MRGPRX2 Activation Causes Increased Skin Reactivity in Patients with Chronic Spontaneous Urticaria. *J Invest Dermatol*. 2021;141(3):678-68
31. Kühn H, Kolkhir P, Babina M, et al. Mas-related G protein-coupled receptor X2 and its activators in dermatologic allergies. *J Allergy Clin Immunol*. 2021;147(2):456-469
32. Youngblood BA, Leung J, Falahati R, et al. Discovery, Function, and Therapeutic Targeting of Siglec-8. *Cell*. 2020 D;10(1):19
33. Zhang S, Cherwinski H, Sedgwick JD, Phillips JH. Molecular mechanisms of CD200 inhibition of mast cell activation. *J Immunol*. 2004;173(11):6786-93
34. Mostmans Y, De Smedt K, Richert B, et al. Markers for the involvement of endothelial cells and the coagulation system in chronic urticaria: A systematic review. *Allergy*. 2021 Mar 25. doi: 10.1111/all.14828
35. Boyman O, Kaegi C, Akdis M, et al. EAACI IG Biologicals task force paper on the use of biologic agents in allergic disorders. *Allergy*. 2015;70(7):727-754
36. Davies AM, Allan EG, Keeble AH, et al. Allosteric mechanism of action of the therapeutic anti-IgE antibody omalizumab. *J Biol Chem*. 2017;292(24):9975-9987
37. Hollis K, Proctor C, McBride D, et al. Comparison of Urticaria Activity Score Over 7 Days (UAS7) Values Obtained from Once-Daily and Twice-Daily Versions: Results from the ASSURE-CSU Study. *Am J Clin Dermatol*. 2018;19(2):267-274
38. Hawro T, Ohanyan T, Schoepke N, et al. The Urticaria Activity Score-Validity, Reliability, and Responsiveness. *J Allergy Clin Immunol Pract*. 2018;6(4):1185-1190
39. Stull D, McBride D, Tian H, et al. Analysis of disease activity categories in chronic spontaneous/idiopathic urticaria. *Br J Dermatol*. 2017;177(4):1093-1101.
40. Mathias SD, Crosby RD, Ros∕@n KE, Zazzali JL. The minimal important difference for measures of urticaria disease activity: Updated findings. *Allergy Asthma Proc*. 2015;36(5):394-398

41. Mathias SD, Crosby RD, Zazzali JL, Maurer M, Saini SS. Evaluating the minimally important difference of the urticaria activity score and other measures of disease activity in patients with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol.* 2012;108(1):20-24.
42. Hawro T, Ohanyan T, Schoepke N, et al. Comparison and interpretability of the available urticaria activity scores. *Allergy* 2018; 73; 251-255
43. Shikhar R, Harding G, Leahy M, Lennox RD. Minimal important difference (MID) of the Dermatology Life Quality Index (DLQI): results from patients with chronic idiopathic urticaria. *Health Qual Life Outcomes.* 2005;3:36.
44. Agache I, Rocha C, Pereira A, et al. Efficacy and safety of treatment with omalizumab for chronic spontaneous urticaria: A systematic review for the EAACI Biologicals Guidelines. *Allergy.* 2021;76(1):59-70
45. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.* 2008;336(7650):924
46. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol.* 2011;64(4):383-94
47. Santesso N, Glenton C, Dahm P, Garner P, Akl EA, Alper B, GRADE Working Group, et al. GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions. *J Clin Epidemiol.* 2020; 119:126-135
48. https://apps.who.int/iris/bitstream/handle/10665/75146/9789241548441_eng.pdf;jsessionid=CA74A1F992AE5574F7B899567C721BC1?sequence=1; accessed on 05.05.2021
49. Maurer M, Altrichter S, Bieber T, et al. Efficacy and safety of omalizumab in patients with chronic urticaria who exhibit IgE against thyroperoxidase. *J Allergy Clin Immunol.* 2011;128(1):202-209
50. Maurer M, Rosén K, Hsieh HJ, et al. Omalizumab for the treatment of chronic idiopathic or spontaneous urticaria [published correction appears in *N Engl J Med.* 2013;368(24):2340-1]. *N Engl J Med.* 2013;368(10):924-935
51. Saini S, Rosen KE, Hsieh HJ, et al. A randomized, placebo-controlled, dose-ranging study of single-dose omalizumab in patients with H1-antihistamine-refractory chronic idiopathic urticaria. *J Allergy Clin Immunol.* 2011;128(3):567-73
52. Saini SS, Bindslev-Jensen C, Maurer M, et al. Efficacy and safety of omalizumab in patients with chronic idiopathic/spontaneous urticaria who remain symptomatic on H1 antihistamines: a randomized, placebo-controlled study [published correction appears in *J Invest Dermatol.* 2015;135(3):925]. *J Invest Dermatol.* 2015;135(1):67-75
53. Staubach P, Metz M, Chapman-Rothe N, et al. Effect of omalizumab on angioedema in H1 -antihistamine-resistant chronic spontaneous urticaria patients: results from X-ACT, a randomized controlled trial [published correction appears in *Allergy.* 2017;72 (9):1430]. *Allergy.* 2016;71(8):1135-1144
54. Metz M, Staubach P, Bauer A, et al. Clinical efficacy of omalizumab in chronic spontaneous urticaria is associated with a reduction of Fc ϵ RI-positive cells in the skin. *Theranostics.* 2017;7(5):1266-1276
55. Jörg L, Pecaric-Petkovic T, Reichenbach S, et al. Double-blind placebo-controlled trial of the effect of omalizumab on basophils in chronic urticaria patients. *Clin Exp Allergy.* 2018;48(2):196-204
56. Hide M, Park HS, Igarashi A, et al. Efficacy and safety of omalizumab in Japanese and Korean patients with refractory chronic spontaneous urticaria. *J Dermatol Sci.* 2017;87(1):70-78
57. Maurer M, Kaplan A, Rosén K, et al. The XTEND-CIU study: Long-term use of omalizumab in chronic idiopathic urticaria. *J Allergy Clin Immunol.* 2018;141(3):1138-1139.
58. Casale TB, Murphy TR, Holden M, Rajput Y, Yoo B, Bernstein JA. Impact of omalizumab on patient-reported outcomes in chronic idiopathic urticaria: Results from a randomized study (XTEND-CIU). *J Allergy Clin Immunol Pract.* 2019;7(7):2487-2490
59. Kaplan A, Ledford D, Ashby M, et al. Omalizumab in patients with symptomatic chronic idiopathic/spontaneous urticaria despite standard combination therapy. *J Allergy Clin Immunol.* 2013;132(1):101-109

60. Deza G, Bertolin-Colilla M, Sanchez S, et al. Basophil FcεRI expression is linked to time to omalizumab response in chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2018;141:2313–2316.
61. Weller K, Ohanyan T, Hawro T, et al. Total IgE levels are linked to the response of chronic spontaneous urticaria patients to omalizumab. *Allergy*. 2018;73:2406–2408
62. Ertas R, Ozyurt K, Atasoy M, Hawro T, Maurer M. The clinical response to omalizumab in chronic spontaneous urticaria patients is linked to and predicted by IgE levels and their change. *Allergy*. 2018;73:705–712
63. de Montjoye L, Darrigade AS, Gimenez-Arnau A, et al. Correlations between disease activity, autoimmunity and biological parameters in patients with chronic spontaneous urticaria. *Eur Ann Allergy Clin Immunol*. 2020
64. Altrichter S, Fok JS, Jiao Q, et al. Total IgE as a Marker for Chronic Spontaneous Urticaria. *Allergy Asthma Immunol Res*. 2021;13(2):206-218
65. Weller K, Groffik A, Church MK, et al. Development and validation of the urticaria control test: a patient-reported outcome instrument for assessing urticaria control. *J Allergy Clin Immunol*. 2014;133(5):1365–1372
66. Weller K, Church MK, Metz M, et al. The response to treatment in chronic spontaneous urticaria depends on how it is measured. *J Allergy Clin Immunol Pract*. 2019;7(6):2055-2056
67. Metz M, Vadasz Z, Kocatürk E, Giménez-Arnau AM. Omalizumab Updosing in Chronic Spontaneous Urticaria: an Overview of Real-World Evidence. *Clin Rev Allergy Immunol*. 2020;59(1):38-45.
68. Khan DA. Hypersensitivity and immunologic reactions to biologics: opportunities for the allergist. *Ann Allergy Asthma Immunol*. 2016;117(2):115-20.4
69. Limb SL, Starke PR, Lee CE, Chowdhury BA. Delayed onset and protracted progression of anaphylaxis after omalizumab administration in patients with asthma. *J Allergy Clin Immunol*. 2007 Dec; 120(6):1378-81
70. Price KS, Hamilton RG. Anaphylactoid reactions in two patients after omalizumab administration after successful long-term therapy. *Allergy Asthma Proc*. 2007;28(3):313-9
71. Macglashan DW Jr, Saini SS. Omalizumab increases the intrinsic sensitivity of human basophils to IgE-mediated stimulation. *J Allergy Clin Immunol*. 2013;132(4):906-11
72. Balbino B, Herviou P, Godon O, et al. The anti-IgE mAb omalizumab induces adverse reactions by engaging Fcγ receptors. *J Clin Invest*. 2020;130(3):1330-1335
73. Bergmann KC, Maurer M, Church MK, Zuberbier T. Anaphylaxis to Mepolizumab and Omalizumab in a Single Patient: Is Polysorbate the Culprit? *J Investig Allergol Clin Immunol*. 2020;30(4):285-287
74. Lieberman PL, Umetsu DT, Carrigan GJ, Rahmaoui A. Anaphylactic reactions associated with omalizumab administration: Analysis of a case-control study. *J Allergy Clin Immunol*. 2016;138(3):913-915
75. Baker DL, Nakamura GR, Lowman HB, Fischer SK. Evaluation of IgE Antibodies to Omalizumab (Xolair®) and Their Potential Correlation to Anaphylaxis. *AAPS J*. 2016;18(1):115-23.
76. Ferastraoaru D, Bax HJ, Bergmann C, et al. AllergoOncology: ultra-low IgE, a potential novel biomarker in cancer—a Position Paper of the European Academy of Allergy and Clinical Immunology (EAACI). *Clin Transl Allergy*. 2020 ;10:32
77. Long A, Rahmaoui A, Rothman KJ, et al. Incidence of malignancy in patients with moderate-to-severe asthma treated with or without omalizumab. *J Allergy Clin Immunol*. 2014; 134: 560-567
78. Johnston A, Smith C, Zheng C, et al. Influence of prolonged treatment with omalizumab on the development of solid epithelial cancer in patients with atopic asthma and chronic idiopathic urticaria: A systematic review and meta-analysis. *Clin Exp Allergy*. 2019;49(10):1291-1305
79. Iribarren C, Rahmaoui A, Long AA, et al. Cardiovascular and cerebrovascular events among patients receiving omalizumab: Results from EXCELS, a prospective cohort study in moderate to severe asthma. *J Allergy Clin Immunol*. 2017;139(5):1489-1495

80. Gill MA, Liu AH, Calatroni A, et al. Enhanced plasmacytoid dendritic cell antiviral responses after omalizumab. *J Allergy Clin Immunol* 2018; 141: 1735– 1743
81. Teach SJ, Gill MA, Togias A, et al. Preseasonal treatment with either omalizumab or an inhaled corticosteroid boost to prevent fall asthma exacerbations. *J Allergy Clin Immunol*. 2015;136(6):1476-1485
82. Tomomatsu K, Oguma T, Baba T, et al; Japan ABPM Research Program. Effectiveness and Safety of Omalizumab in Patients with Allergic Bronchopulmonary Aspergillosis Complicated by Chronic Bacterial Infection in the Airways. *Int Arch Allergy Immunol*. 2020;181(7):499-506
83. Cruz AA, Lima F, Sarinho E, et al. Safety of anti-immunoglobulin E therapy with omalizumab in allergic patients at risk of geohelminth infection. *Clin Exp Allergy*. 2007;37(2):197-207
84. Winthrop KL, Mariette X, Silva JT, et al. ESCMID Study Group for Infections in Compromised Hosts (ESGICH) Consensus Document on the safety of targeted and biological therapies: an infectious diseases perspective (Soluble immune effector molecules [II]: agents targeting interleukins, immunoglobulins and complement factors). *Clin Microbiol Infect*. 2018;24 Suppl 2:S21-S40.
85. Gasser P, Tarchevskaya SS, Guntern P, et al. The mechanistic and functional profile of the therapeutic anti-IgE antibody ligelizumab differs from omalizumab. *Nat Commun*. 2020;11(1):165
86. Maurer M, Giménez-Arnau AM, Sussman G, et al. Ligelizumab for chronic spontaneous urticaria. *N Engl J Med*. 2019;381:1321–1332
87. Brightbill HD, Lin YL, Lin Z et al. Quilizumab is an afucosylated humanized anti-M1 prime therapeutic antibody. *Clin Antinflamm AntiAllergy Drugs*. 2014; 1: 24-31
88. Harris JM, Cabanski CR, Scheerens H, et al. A randomized trial of quilizumab in adults with refractory chronic spontaneous urticaria. *J Allergy Clin Immunol*. 2016;138:1730–1732
89. Harris JM, Maciucă R, Bradley MS, et al. A randomized trial of the efficacy and safety of quilizumab in adults with inadequately controlled allergic asthma. *Respir Res*. 2016;17:29.
90. http://gi-innovation.com/en/images/content/GI-301_PDF.pdf; accessed 07.05.2021
91. Lee JK, Simpson RS. Dupilumab as a novel therapy for difficult to treat chronic spontaneous urticaria. *J Allergy Clin Immunol Pract*. 2019; 7(5):1659-1661
92. Magerl M, Terhorst D, Metz M, et al. Benefit of mepolizumab treatment in a patient with chronic spontaneous urticaria. *J Dtsch Dermatol Ges*. 2018; 16(4):477-478.
93. Maurer M, Altrichter S, Metz M, et al. Benefit from reslizumab treatment in a patient with chronic spontaneous urticaria and cold urticaria. *J Eur Acad Dermatol Venereol*. 2018; 32(3):e112-e113
94. Bernstein JA, Singh U, Rao MB, et al. Benralizumab for Chronic Spontaneous Urticaria. *N Engl J Med*. 2020; 383(14):1389-1391
95. Bernstein JA, Singh U, Rao MB, et al. Treatment of chronic spontaneous urticaria with benralizumab: Report of primary endpoint per-protocol analysis and exploratory endpoints. *Allergy*. 2021;76(4):1277-1280
96. Altrichter S, Staubach P, Pasha M et al. Efficacy and safety data of AK002, an anti-siglec-8 monoclonal antibody, in patients with multiple forms of uncontrolled chronic urticaria (CU): Results from an open-label phase 2a study. *Allergy*. 2020; 74: 120
97. Dellon ES, Peterson KA, Murray JA et al. Anti-Siglec-8 antibody for eosinophilic gastritis and duodenitis. *N Engl J Med*. 2020; 383: 1624–1634
98. Maurer M, Crew L, Murphy M et al. CDX-0159, an anti-KIT monoclonal antibody, demonstrates dose-dependent reductions in serum tryptase and a favorable safety profile in a phase 1a healthy volunteer study. *Allergy*. 2020; 75:280.
99. Bernstein JA, Lang DM, Khan DA, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. *J Allergy Clin Immunol*. 2014;133(5):1270-7
100. Caffarelli C, Paravati F, El Hachem M, et al. Management of chronic urticaria in children: a clinical guideline. *Ital J Pediatr*. 2019;45(1):101.

101. Choi JH, Lee DH, Song WJ, et al. The KAAACI/KDA Evidence-Based Practice Guidelines for Chronic Spontaneous Urticaria in Korean Adults and Children: Part 2. Management of H1-Antihistamine-Refractory Chronic Urticaria. *Allergy Asthma Immunol Res.* 2020;12(5):750-770
102. Metz M, Altrichter S, Buttgereit T, et al. The Diagnostic Workup in Chronic Spontaneous Urticaria-What to Test and Why. *J Allergy Clin Immunol Pract.* 2021:S2213-2198(21)00435-9
103. Fok JS, Kolkhir P, Church MK, Maurer M. Predictors of treatment response in chronic spontaneous urticaria. *Allergy.* 2021 Feb 4. doi: 10.1111/all.14757. Epub ahead of print
104. Breiteneder H, Peng YQ, Agache I, et al. Biomarkers for diagnosis and prediction of therapy responses in allergic diseases and asthma. *Allergy.* 2020;75(12):3039-3068
105. Chan MA, Gigliotti NM, Dotson AL, Rosenwasser LJ. Omalizumab may decrease IgE synthesis by targeting membrane IgE+ human B cells. *Clin Transl Allergy.* 2013; 3(1):29
106. Eyerich S, Metz M, Bossios A, Eyerich K. New biological treatments for asthma and skin allergies. *Allergy.* 2020 Mar;75(3):546-560
107. Metz M, Maurer M. Use of biologics in chronic spontaneous urticaria - beyond omalizumab therapy? *Allergol Select.* 2021;5:89-95.
108. Mansur AH, Srivastava S, Mitchell V, Sullivan J, Kasujee I. Long-term clinical outcomes of omalizumab therapy in severe allergic asthma: Study of efficacy and safety. *Respir Med.* 2017;124:36-43
109. Nieto García A, Garriga-Baraut T, Plaza Martín AM, et al. Omalizumab outcomes for up to 6 years in pediatric patients with severe persistent allergic asthma. *Pediatr Allergy Immunol.* 2021 Feb 22. doi: 10.1111/pai.13484. Epub ahead of print.
110. Nakamura N, Kashitani Y, Yoshisue H, Nagasaki M, Sasajima T. Real-life long-term safety and effectiveness of omalizumab in Japanese pediatric patients with severe allergic asthma: A post-marketing surveillance. *Allergol Int.* 2021:S1323-8930(21)00001-0
111. Kanters TA, Thio HB, Hakkaart L. Cost-effectiveness of omalizumab for the treatment of chronic spontaneous urticaria. *Br J Dermatol.* 2018;179(3):702-708
112. Graham J, McBride D, Stull D, et al. Cost Utility of Omalizumab Compared with Standard of Care for the Treatment of Chronic Spontaneous Urticaria. *Pharmacoeconomics.* 2016;34(8):815-27